

Examples of Different Results using car package

Kyun-Seop Bae MD PhD

2023-08-29

Contents

| | | |
|----------|---|-----------|
| 1 | Tested Version and Books used for the Validation | 3 |
| 1.1 | Packages Used | 3 |
| 1.2 | Books and Articles used for the Test | 3 |
| 2 | Snee EMS ANOVA 1974 | 4 |
| 3 | Goodnight | 5 |
| 3.1 | p33 | 5 |
| 4 | SAS for Linear Models 4e | 6 |
| 4.1 | p403 | 6 |
| 4.2 | p417 | 7 |
| 4.3 | p431 | 8 |
| 5 | Sahai - Unbalanced | 9 |
| 5.1 | Table 15.3 | 9 |
| 5.2 | Table 16.3 | 10 |
| 6 | Federer - Variations | 11 |
| 6.1 | Example 2.2 | 11 |
| 6.2 | Example 3.1 | 12 |
| 6.3 | Appendix 3.1 p94 | 13 |
| 6.4 | Example 5.1 | 14 |
| 6.5 | Example 7.1 | 16 |
| 6.6 | Example 7.3 | 17 |
| 6.7 | Example 8.1 | 18 |
| 6.8 | Example 9.2 | 19 |
| 6.9 | Example 10.1 | 20 |
| 7 | Hinkelmann & Kempthorne - Volume 1 | 22 |
| 7.1 | p410 | 22 |
| 8 | Searle - Linear Models 2e | 23 |
| 8.1 | 7.2 (p390, 59%) | 23 |
| 8.2 | 7.2 (p393, 60%) | 24 |

| | | |
|-----------|---|-----------|
| 9 | Web site examples | 25 |
| 9.1 | https://github.com/djnavarro/psyr | 25 |
| 10 | Bioequivalence (BE) data example | 27 |
| 11 | Sesssion Information | 29 |

1 Tested Version and Books used for the Validation

1.1 Packages Used

- ‘sasLM’ version: 0.9.12
- ‘SAS’ version: 9.4 Licensed and University Edition
- ‘car’ version: 3.1.2
- R version: R version 4.3.1 (2023-06-16 ucrt)

The ‘car’ package is not necessary for ‘sasLM.’ It is used for the comparison of the results.

If you see any difference between ‘car’ and ‘sasLM’, ‘SAS’ results coincide with ‘sasLM’, not with ‘car’.

Before ‘sasLM’ is available on CRAN, you can download using the following command in R.

```
install.packages("sasLM", repos="http://r.acr.kr")
```

1.2 Books and Articles used for the Test

1. Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. *J Qual Tech.* 1974;6(3):128-137.
2. Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User’s Group, SAS Institute, Raleigh, N.C. 1976.
3. Littell RC, Stroup WW, Freund RJ. *SAS for Linear Models 4e*. John Wiley & Sons Inc. 2002.
4. Sahai H, Ojeda MM. *Analysis of Variance for Random Models Volume 2 Unbalanced Data*. 2005.
5. Federer WT, King F. *Variations on Split Plot and Split Block Experiment Designs*. John Wiley & Sons Inc. 2007.
6. Hinkelmann K, Kempthorne O. *Design and Analysis of Experiments Volume 1 Introduction to Experimental Design*. 2e. John Wiley & Sons Inc. 2008.
7. Searle SR, Gruber MHJ. *Linear Models 2e*, Kindle Edition. John Wiley & Sons Inc. 2016.

2 Snee EMS ANOVA 1974

Reference

- Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.

(1) MODEL

```
Snee = read.csv("http://r.acr.kr/Snee_EMS_ANOVA1974.csv")
Snee = af(Snee, c("Machine", "Analyst", "Test", "Day"))
aov3(Y ~ Day/Machine/Analyst/Test, Snee)
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|--------------------------|-----|--------|---------|---------|--------|
| MODEL | 167 | 751.27 | 4.4986 | | |
| Day | 41 | 359.44 | 8.7669 | | |
| Day:Machine | 42 | 199.40 | 4.7477 | | |
| Day:Machine:Analyst | 42 | 118.80 | 2.8285 | | |
| Day:Machine:Analyst:Test | 42 | 70.30 | 1.6739 | | |
| RESIDUALS | 0 | 0.00 | | | |
| CORRECTED TOTAL | 167 | 751.27 | | | |

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Day/Machine/Analyst/Test, Snee), type=3, singular.ok=TRUE)
# NOT WORKING
```

3 Goodnight

Reference

- Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User's Group, SAS Institute, Raleigh, N.C. 1976.

3.1 p33

(2) MODEL

```
p33 = read.csv("http://r.acr.kr/Goodnight-p33.csv")
p33 = af(p33, c("A", "B"))
aov3(y ~ A + B + A:B, p33) # p35
```

Response : y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|--------|---------|---------|--------|
| MODEL | 4 | 34.905 | 8.7261 | | |
| A | 1 | 3.028 | 3.0276 | | |
| B | 1 | 23.523 | 23.5225 | | |
| A:B | 1 | 0.008 | 0.0081 | | |
| RESIDUALS | 0 | 0.000 | | | |
| CORRECTED TOTAL | 4 | 34.905 | | | |

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(y ~ A + B + A:B, p33), type=3, singular.ok=TRUE) # NOT WORKING
```

4 SAS for Linear Models 4e

Reference

- Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.

4.1 p403

(3) MODEL

```
p403 = read.table("http://r.acr.kr/sas4lm/p403.txt", header=TRUE)
p403 = af(p403, c("PATIENT", "VISIT"))
aov3(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT, p403)
```

Response : HR

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|------------------|----|--------|---------|---------|---------------|
| MODEL | 29 | 6408.7 | 220.989 | 3.9120 | 3.127e-05 *** |
| SEQUENCE | 5 | 701.2 | 140.237 | 2.4825 | 0.04665 * |
| VISIT | 2 | 146.8 | 73.389 | 1.2991 | 0.28350 |
| DRUG | 2 | 344.0 | 171.975 | 3.0443 | 0.05826 . |
| RESIDS | 1 | 309.2 | 309.174 | 5.4731 | 0.02414 * |
| RESIDT | 1 | 0.8 | 0.840 | 0.0149 | 0.90351 |
| SEQUENCE:PATIENT | 18 | 4692.3 | 260.685 | 4.6147 | 2.210e-05 *** |
| RESIDUALS | 42 | 2372.6 | 56.490 | | |
| CORRECTED TOTAL | 71 | 8781.3 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT,
         p403), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: HR

| | Sum Sq | Df | F values | Pr(>F) |
|------------------|--------|----|----------|--------------|
| SEQUENCE | 0.0 | 0 | | |
| VISIT | 146.8 | 2 | 1.2991 | 0.28350 |
| DRUG | 343.9 | 2 | 3.0443 | 0.05826 . |
| RESIDS | 309.2 | 1 | 5.4731 | 0.02414 * |
| RESIDT | 0.8 | 1 | 0.0149 | 0.90351 |
| SEQUENCE:PATIENT | 4692.3 | 18 | 4.6147 | 2.21e-05 *** |
| Residuals | 2372.6 | 42 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

4.2 p417

(4) MODEL

```
p417 = read.table("http://r.acr.kr/sas4lm/p417.txt", header=TRUE)
p417 = af(p417, c("TRT", "POT", "PLANT"))
aov3(Y ~ TRT + POT %in% TRT, p417) # p418 Output 11.28
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|---------|---------|---------|---------------|
| MODEL | 7 | 267.226 | 38.175 | 12.433 | 7.522e-05 *** |
| TRT | 2 | 200.111 | 100.055 | 32.586 | 8.626e-06 *** |
| TRT:POT | 5 | 30.306 | 6.061 | 1.974 | 0.1499 |
| RESIDUALS | 13 | 39.917 | 3.071 | | |
| CORRECTED TOTAL | 20 | 307.143 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ TRT + POT %in% TRT, p417), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|--------|----|----------|-----------|
| TRT | 22.310 | 1 | 7.266 | 0.01835 * |
| TRT:POT | 30.306 | 5 | 1.974 | 0.14991 |
| Residuals | 39.917 | 13 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

4.3 p431

(5) MODEL

```
p431 = read.table("http://r.acr.kr/sas4lm/p431.txt", header=TRUE)
p431 = af(p431, c("line", "sire", "agedam", "steerno"))
aov3(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431)
```

Response : avdlygn

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|--------|---------|---------|-------------|
| MODEL | 16 | 2.5275 | 0.15797 | 3.1437 | 0.001091 ** |
| line | 2 | 0.1362 | 0.06810 | 1.3553 | 0.267560 |
| agedam | 2 | 0.1301 | 0.06505 | 1.2946 | 0.283392 |
| age | 1 | 0.3813 | 0.38128 | 7.5878 | 0.008277 ** |
| intlw | 1 | 0.2697 | 0.26970 | 5.3674 | 0.024830 * |
| line:sire | 6 | 0.9739 | 0.16231 | 3.2303 | 0.009543 ** |
| line:agedam | 4 | 0.4534 | 0.11336 | 2.2560 | 0.076821 . |
| RESIDUALS | 48 | 2.4119 | 0.05025 | | |
| CORRECTED TOTAL | 64 | 4.9394 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

p433 Output 11.40

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431),
      type=3, singular.ok=TRUE) # NOT OK for line
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: avdlygn

| | Sum Sq | Df | F values | Pr(>F) |
|-------------|---------|----|----------|-------------|
| line | 0.00000 | 0 | | |
| agedam | 0.13011 | 2 | 1.2946 | 0.283392 |
| age | 0.38128 | 1 | 7.5878 | 0.008277 ** |
| intlw | 0.26970 | 1 | 5.3674 | 0.024830 * |
| line:sire | 0.97389 | 6 | 3.2303 | 0.009543 ** |
| line:agedam | 0.45343 | 4 | 2.2560 | 0.076821 . |
| Residuals | 2.41192 | 48 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5 Sahai - Unbalanced

Reference

- Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.

5.1 Table 15.3

(6) MODEL

```
T15.3 = read.table("http://r.acr.kr/sahai/T15.3.txt")
colnames(T15.3) = c("Dam", "Sire", "pH")
T15.3 = af(T15.3, c("Dam", "Sire"))
aov3(pH ~ Dam/Sire, T15.3) # p301
```

Response : pH

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|---------|-----------|---------|---------------|
| MODEL | 36 | 0.25804 | 0.0071678 | 2.8977 | 7.200e-06 *** |
| Dam | 14 | 0.17940 | 0.0128146 | 5.1805 | 1.347e-07 *** |
| Dam:Sire | 22 | 0.08002 | 0.0036374 | 1.4705 | 0.09662 . |
| RESIDUALS | 123 | 0.30425 | 0.0024736 | | |
| CORRECTED TOTAL | 159 | 0.56229 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(pH ~ Dam/Sire, T15.3), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: pH

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|----------|-----|----------|---------------|
| Dam | 0.081011 | 6 | 5.4584 | 4.898e-05 *** |
| Dam:Sire | 0.080024 | 22 | 1.4705 | 0.09662 . |
| Residuals | 0.304253 | 123 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.2 Table 16.3

(7) MODEL

```
T16.3 = read.csv("http://r.acr.kr/sahai/T16.3.csv")
colnames(T16.3) = c("Plot", "Sample", "Subsample", "Residue")
T16.3 = af(T16.3, c("Plot", "Sample", "Subsample"))
aov3(Residue ~ Plot/Sample/Subsample, T16.3) # p344
```

```
Response : Residue
             Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          54 3.1897 0.059069  5.8842 1.476e-05 ***
Plot           10 1.7869 0.178686 17.7998 2.547e-08 ***
Plot:Sample    22 0.9917 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.3576 0.016253  1.6191 0.1330632
RESIDUALS      22 0.2208 0.010039
CORRECTED TOTAL 76 3.4106
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(Residue ~ Plot/Sample/Subsample, T16.3), type=3, singular.ok=TRUE)
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```
Response: Residue
             Sum Sq Df F values   Pr(>F)
Plot          0.00000  0
Plot:Sample   0.36613 11  3.3156 0.00805 **
Plot:Sample:Subsample 0.35758 22  1.6191 0.13306
Residuals     0.22085 22
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
# NOT OK
```

6 Federer - Variations

Reference

- Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.

6.1 Example 2.2

(8) MODEL

```
ex2.2 = read.table("http://r.acr.kr/split/sbex2_2.txt", header=TRUE)
ex2.2 = af(ex2.2, c("Row", "Column", "R", "S"))
aov3(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2)
```

```
Response : Y
              Df  Sum Sq Mean Sq F value Pr(>F)
MODEL          99 22310.4 225.36
Row             0
R               4   1159.8 289.94
S               3    351.9 117.29
R:S              12   826.0  68.83
Row:R            0
S:Column         12   3863.3 321.94
R:S:Column       48 11982.3 249.63
RESIDUALS        0     0.0
CORRECTED TOTAL 99 22310.4
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2), type=3,
      singular.ok=TRUE) # NOT WORKING
```

6.2 Example 3.1

(9) MODEL

```
ex3.1a = read.table("http://r.acr.kr/split/Ex3.1-example.txt", header=TRUE)
ex3.1a = af(ex3.1a, c("row", "P", "column", "R", "S"))
aov3(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
      S:R:P + R:S:P:row, ex3.1a)
```

Response : height

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|--------|---------|---------|--------|
| MODEL | 199 | 7534.8 | 37.86 | | |
| row | 4 | 2017.0 | 504.26 | | |
| R | 4 | 90.6 | 22.66 | | |
| P | 1 | 253.1 | 253.13 | | |
| S | 3 | 16.4 | 5.46 | | |
| R:S | 12 | 195.0 | 16.25 | | |
| row:P | 4 | 167.2 | 41.81 | | |
| R:P | 4 | 505.0 | 126.24 | | |
| P:S | 3 | 14.3 | 4.77 | | |
| row:R:P | 32 | 2933.5 | 91.67 | | |
| row:P:S | 24 | 234.7 | 9.78 | | |
| R:P:S | 12 | 100.3 | 8.36 | | |
| row:R:P:S | 96 | 1007.5 | 10.49 | | |
| RESIDUALS | 0 | 0.0 | | | |
| CORRECTED TOTAL | 199 | 7534.8 | | | |

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
          S:P:row + S:R:P + R:S:P:row, ex3.1a), type=3, singular.ok=TRUE)
# NOT WORKING
```

6.3 Appendix 3.1 p94

(10) MODEL

```
ex3.1b = read.table("http://r.acr.kr/split/spexvar3.txt", header=TRUE)
ex3.1b = af(ex3.1b, c("rep", "var", "nit", "row", "col"))
aov3(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b)
```

Response : yield

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|--------|---------|---------|---------------|
| MODEL | 37 | 48090 | 1299.7 | 11.3414 | 6.734e-11 *** |
| rep | 2 | 5943 | 2971.3 | 25.9273 | 1.449e-07 *** |
| var | 2 | 2800 | 1399.9 | 12.2155 | 0.0001005 *** |
| nit | 3 | 11978 | 3992.6 | 34.8397 | 1.775e-10 *** |
| row | 9 | 945 | 105.0 | 0.9162 | 0.5230151 |
| col | 2 | 3171 | 1585.7 | 13.8373 | 4.012e-05 *** |
| rep:var | 4 | 998 | 249.4 | 2.1767 | 0.0926008 . |
| var:nit | 6 | 478 | 79.6 | 0.6949 | 0.6553307 |
| RESIDUALS | 34 | 3896 | 114.6 | | |
| CORRECTED TOTAL | 71 | 51986 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b),
      type=3, singular.ok=TRUE) # NOT OK for var
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: yield

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|---------|----|----------|---------------|
| rep | 5942.5 | 2 | 25.9273 | 1.449e-07 *** |
| var | 0.0 | 0 | | |
| nit | 11977.9 | 3 | 34.8397 | 1.775e-10 *** |
| row | 945.0 | 9 | 0.9162 | 0.5230 |
| col | 3171.5 | 2 | 13.8373 | 4.012e-05 *** |
| rep:var | 997.8 | 4 | 2.1767 | 0.0926 . |
| var:nit | 477.8 | 6 | 0.6949 | 0.6553 |
| Residuals | 3896.4 | 34 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.4 Example 5.1

(11) MODEL

```
ex5.1 = read.table("http://r.acr.kr/split/sbsp.txt", header=TRUE)
ex5.1 = af(ex5.1, c("R", "A", "C", "B", "Tx"))
aov3(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|---------|---------|---------|---------------|
| MODEL | 24 | 196.238 | 8.1766 | 7.0476 | 0.0008758 *** |
| R | 2 | 22.186 | 11.0928 | 9.5611 | 0.0039244 ** |
| A | 1 | 15.185 | 15.1853 | 13.0886 | 0.0040418 ** |
| C | 2 | 1.010 | 0.5049 | 0.4352 | 0.6578395 |
| B | 1 | 1.792 | 1.7922 | 1.5448 | 0.2397515 |
| Tx | 5 | 103.333 | 20.6667 | 17.8131 | 6.055e-05 *** |
| R:A | 2 | 27.426 | 13.7132 | 11.8197 | 0.0018198 ** |
| C:B | 2 | 0.085 | 0.0424 | 0.0366 | 0.9642020 |
| A:Tx | 4 | 2.655 | 0.6636 | 0.5720 | 0.6886524 |
| B:Tx | 4 | 2.050 | 0.5126 | 0.4418 | 0.7761730 |
| RESIDUALS | 11 | 12.762 | 1.1602 | | |
| CORRECTED TOTAL | 35 | 209.000 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|---------|----|----------|---------------|
| R | 22.186 | 2 | 9.5611 | 0.003924 ** |
| A | 0.000 | 0 | | |
| C | 1.010 | 2 | 0.4352 | 0.657839 |
| B | 0.000 | 0 | | |
| Tx | 103.333 | 5 | 17.8131 | 6.055e-05 *** |
| R:A | 27.426 | 2 | 11.8197 | 0.001820 ** |
| C:B | 0.085 | 2 | 0.0366 | 0.964202 |
| A:Tx | 2.655 | 4 | 0.5720 | 0.688652 |
| B:Tx | 2.050 | 4 | 0.4418 | 0.776173 |
| Residuals | 12.762 | 11 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(12) MODEL

```
aov3(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

```
Response : Y
            Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        28 204.200  7.2929 10.6354 0.0017194 ***
R             2   28.112 14.0562 20.4986 0.0011846 ***
A             1   14.655 14.6551 21.3720 0.0024176 ***
C             2    0.471  0.2356  0.3436 0.7205632
B             1    1.769  1.7694  2.5804 0.1522328
Tx            5 103.815 20.7630 30.2793 0.0001336 ***
R:A           1    2.017  2.0174  2.9420 0.1300172
C:B           1    0.644  0.6445  0.9399 0.3646045
A:Tx          4    2.951  0.7378  1.0760 0.4358837
B:Tx          4    3.553  0.8882  1.2954 0.3579988
A:B:Tx        4    7.962  1.9905  2.9029 0.1038803
RESIDUALS     7    4.800  0.6857
CORRECTED TOTAL 35 209.000
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```
Response: Y
            Sum Sq Df F values    Pr(>F)
R         11.643  1 16.9793 0.004456 ***
A         0.000  0
C         0.002  1  0.0025 0.961483
B         0.000  0
Tx        89.178  3 43.3503 6.87e-05 ***
R:A       2.017  1  2.9420 0.130017
C:B       0.644  1  0.9399 0.364604
A:Tx      0.543  3  0.2640 0.849381
B:Tx      3.384  3  1.6451 0.264128
A:B:Tx    7.962  4  2.9029 0.103880
Residuals 4.800  7
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

6.5 Example 7.1

(13) MODEL

```
ex7.1 = read.table("http://r.acr.kr/split/asped.txt", header=TRUE)
ex7.1 = af(ex7.1, c("R", "G", "F"))
aov3(Y ~ R + G + R:G + F + F:G, ex7.1)
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|--------|---------|---------|---------------|
| MODEL | 95 | 577.83 | 6.0824 | 5.3082 | 1.068e-05 *** |
| R | 3 | 5.75 | 1.9167 | 1.6727 | 0.1994 |
| G | 27 | 343.48 | 12.7216 | 11.1025 | 4.286e-08 *** |
| F | 2 | 50.50 | 25.2525 | 22.0385 | 3.686e-06 *** |
| R:G | 9 | 11.75 | 1.3056 | 1.1394 | 0.3749 |
| G:F | 54 | 77.98 | 1.4441 | 1.2603 | 0.2718 |
| RESIDUALS | 24 | 27.50 | 1.1458 | | |
| CORRECTED TOTAL | 119 | 605.33 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + G + R:G + F + F:G, ex7.1), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|---------|----|----------|---------------|
| R | 0.000 | 0 | | |
| G | 202.417 | 3 | 58.8848 | 3.258e-11 *** |
| F | 50.505 | 2 | 22.0385 | 3.686e-06 *** |
| R:G | 11.750 | 9 | 1.1394 | 0.3749 |
| G:F | 77.983 | 54 | 1.2603 | 0.2718 |
| Residuals | 27.500 | 24 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.6 Example 7.3

(14) MODEL

```
ex7.3 = read.table("http://r.acr.kr/split/assped.txt", header=TRUE)
ex7.3 = af(ex7.3, c("R", "T", "G", "F"))
aov3(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3)
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|--------|---------|----------|---------------|
| MODEL | 155 | 656.12 | 4.233 | 13.4461 | 3.997e-14 *** |
| R | 3 | 12.49 | 4.162 | 13.2206 | 5.655e-06 *** |
| T | 1 | 11.16 | 11.158 | 35.4430 | 8.021e-07 *** |
| G | 22 | 389.01 | 17.682 | 56.1668 | < 2.2e-16 *** |
| F | 2 | 120.56 | 60.282 | 191.4828 | < 2.2e-16 *** |
| R:T | 3 | 1.15 | 0.384 | 1.2206 | 0.316281 |
| T:G | 22 | 18.42 | 0.837 | 2.6601 | 0.004445 ** |
| T:F | 2 | 0.82 | 0.411 | 1.3060 | 0.283432 |
| G:F | 44 | 23.47 | 0.533 | 1.6943 | 0.053191 . |
| R:T:G | 12 | 8.78 | 0.731 | 2.3235 | 0.025315 * |
| T:G:F | 44 | 10.74 | 0.244 | 0.7753 | 0.790640 |
| RESIDUALS | 36 | 11.33 | 0.315 | | |
| CORRECTED TOTAL | 191 | 667.45 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3),
      type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|---------|----|------------------------|--------|
| R | 0.000 | 0 | | |
| T | 0.000 | 0 | | |
| G | 73.444 | 2 | 116.6471 < 2.2e-16 *** | |
| F | 120.563 | 2 | 191.4828 < 2.2e-16 *** | |
| R:T | 0.000 | 0 | | |
| T:G | 5.778 | 2 | 9.1765 0.0006018 *** | |
| T:F | 0.822 | 2 | 1.3060 0.2834316 | |
| G:F | 23.469 | 44 | 1.6943 0.0531910 . | |
| R:T:G | 8.778 | 12 | 2.3235 0.0253153 * | |
| T:G:F | 10.740 | 44 | 0.7753 0.7906401 | |
| Residuals | 11.333 | 36 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.7 Example 8.1

(15) MODEL

```
ex8.1 = read.table("http://r.acr.kr/split/asbed.txt", header=TRUE)
ex8.1 = af(ex8.1, c("R", "A", "B"))
aov3(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1)
```

Response : Y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|--------|---------|---------|--------|
| MODEL | 104 | 3951.8 | 37.999 | | |
| R | 2 | 372.2 | 186.111 | | |
| A | 12 | 572.3 | 47.692 | | |
| B | 8 | 185.8 | 23.231 | | |
| R:A | 6 | 50.0 | 8.333 | | |
| R:B | 4 | 87.4 | 21.861 | | |
| A:B | 60 | 1012.3 | 16.871 | | |
| R:A:B | 12 | 49.0 | 4.083 | | |
| RESIDUALS | 0 | 0.0 | | | |
| CORRECTED TOTAL | 104 | 3951.8 | | | |

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1), type="III",
singular.ok=TRUE) # NOT WORKING
```

6.8 Example 9.2

(16) MODEL

```
ex9.2 = read.table("http://r.acr.kr/split/Ex9.2-sbex.txt", header=TRUE)
ex9.2 = af(ex9.2, c("rep", "hyb", "gen"))
aov3(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2)
```

Response : yield

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|---------|---------|---------|---------------|
| MODEL | 40 | 247.813 | 6.1953 | 4.4606 | 0.0011186 ** |
| rep | 1 | 0.167 | 0.1667 | 0.1200 | 0.7335481 |
| hyb | 9 | 66.796 | 7.4218 | 5.3437 | 0.0018370 ** |
| gen | 2 | 30.671 | 15.3356 | 11.0416 | 0.0009707 *** |
| rep:hyb | 8 | 67.000 | 8.3750 | 6.0300 | 0.0011569 ** |
| rep:gen | 2 | 12.111 | 6.0556 | 4.3600 | 0.0308015 * |
| hyb:gen | 18 | 60.504 | 3.3613 | 2.4201 | 0.0408545 * |
| RESIDUALS | 16 | 22.222 | 1.3889 | | |
| CORRECTED TOTAL | 56 | 270.035 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2), type=3,
singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients

sums of squares computed by model comparison

Anova Table (Type III tests)

Response: yield

| | Sum Sq | Df | F values | Pr(>F) |
|-----------|--------|----|----------|---------------|
| rep | 0.000 | 0 | | |
| hyb | 66.704 | 8 | 6.0033 | 0.0011847 ** |
| gen | 30.671 | 2 | 11.0416 | 0.0009707 *** |
| rep:hyb | 67.000 | 8 | 6.0300 | 0.0011569 ** |
| rep:gen | 12.111 | 2 | 4.3600 | 0.0308015 * |
| hyb:gen | 60.504 | 18 | 2.4201 | 0.0408545 * |
| Residuals | 22.222 | 16 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.9 Example 10.1

(17) MODEL

```
ex10.1 = read.table("http://r.acr.kr/split/Ex10.1-New.txt", header=TRUE)
ex10.1 = af(ex10.1, c("Site", "Block", "A", "B", "C"))
f10.1 = Yield ~ Site/Block + A/Site + B/Site + A:B + A:B:Site + A:B:Site:Block +
         C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site + A:B:C:Site
aov3(f10.1, ex10.1)
```

Response : Yield

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|-----|------------|-----------|------------|-------------|
| MODEL | 239 | 1639561484 | 6860090 | 2.1620e+03 | < 2e-16 *** |
| Site | 3 | 552717 | 184239 | 5.8064e+01 | < 2e-16 *** |
| A | 4 | 1387680917 | 346920229 | 1.0933e+05 | < 2e-16 *** |
| B | 1 | 100939695 | 100939695 | 3.1812e+04 | < 2e-16 *** |
| C | 3 | 19356264 | 6452088 | 2.0334e+03 | < 2e-16 *** |
| Site:Block | 8 | 7062320 | 882790 | 2.7822e+02 | < 2e-16 *** |
| Site:A | 12 | 34068 | 2839 | 8.9470e-01 | 0.55301 |
| Site:B | 3 | 1618 | 539 | 1.6990e-01 | 0.91662 |
| A:B | 4 | 31444008 | 7861002 | 2.4775e+03 | < 2e-16 *** |
| A:C | 12 | 26075792 | 2172983 | 6.8483e+02 | < 2e-16 *** |
| B:C | 3 | 23901387 | 7967129 | 2.5109e+03 | < 2e-16 *** |
| Site:C | 9 | 47625 | 5292 | 1.6677e+00 | 0.09747 . |
| Site:A:B | 12 | 33737 | 2811 | 8.8600e-01 | 0.56185 |
| A:B:C | 12 | 41996729 | 3499727 | 1.1030e+03 | < 2e-16 *** |
| Site:A:C | 36 | 104110 | 2892 | 9.1140e-01 | 0.61768 |
| Site:B:C | 9 | 61111 | 6790 | 2.1400e+00 | 0.02701 * |
| Site:Block:A:B | 72 | 186911 | 2596 | 8.1810e-01 | 0.84155 |
| Site:A:B:C | 36 | 82475 | 2291 | 7.2200e-01 | 0.87941 |
| RESIDUALS | 240 | 761522 | 3173 | | |
| CORRECTED TOTAL | 479 | 1640323006 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(f10.1, ex10.1), type=3, singular.ok=TRUE) # NOT OK for Site:Block
```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

| Response: Yield | Sum Sq | Df | F values | Pr(>F) |
|-----------------|------------|----|------------|-------------|
| Site | 552717 | 3 | 5.8064e+01 | < 2e-16 *** |
| A | 1387680917 | 4 | 1.0933e+05 | < 2e-16 *** |
| B | 100939695 | 1 | 3.1812e+04 | < 2e-16 *** |
| C | 19356264 | 3 | 2.0334e+03 | < 2e-16 *** |
| Site:Block | 0 | 0 | | |

| | | | | | | | | | | | |
|----------------|----------|-------|------------|-------------|------|-----|------|------|-----|-----|---|
| Site:A | 34068 | 12 | 8.9470e-01 | 0.55301 | | | | | | | |
| Site:B | 1618 | 3 | 1.6990e-01 | 0.91662 | | | | | | | |
| A:B | 31444008 | 4 | 2.4775e+03 | < 2e-16 *** | | | | | | | |
| A:C | 26075792 | 12 | 6.8483e+02 | < 2e-16 *** | | | | | | | |
| B:C | 23901388 | 3 | 2.5109e+03 | < 2e-16 *** | | | | | | | |
| Site:C | 47625 | 9 | 1.6677e+00 | 0.09747 . | | | | | | | |
| Site:A:B | 33737 | 12 | 8.8600e-01 | 0.56185 | | | | | | | |
| A:B:C | 41996729 | 12 | 1.1030e+03 | < 2e-16 *** | | | | | | | |
| Site:A:C | 104110 | 36 | 9.1140e-01 | 0.61768 | | | | | | | |
| Site:B:C | 61111 | 9 | 2.1400e+00 | 0.02701 * | | | | | | | |
| Site:Block:A:B | 186911 | 72 | 8.1810e-01 | 0.84155 | | | | | | | |
| Site:A:B:C | 82475 | 36 | 7.2200e-01 | 0.87941 | | | | | | | |
| Residuals | 761522 | 240 | | | | | | | | | |
| | --- | | | | | | | | | | |
| Signif. codes: | 0 | '***' | 0.001 | '**' | 0.01 | '*' | 0.05 | '..' | 0.1 | ' ' | 1 |

7 Hinkelmann & Kempthorne - Volume 1

Reference

- Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.

7.1 p410

(18) MODEL

```
v1p410 = read.table("http://r.acr.kr/kemp/v1p410.txt", head=TRUE)
v1p410$carry = ifelse(v1p410$carry == 0, 3, v1p410$carry)
v1p410 = af(v1p410,c("period", "sequence", "steer", "trt", "carry"))
aov3(y ~ period + sequence + steer:sequence + trt + carry, v1p410) # OK
```

Response : y

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|---------|---------|---------|---------------|
| MODEL | 17 | 1302.51 | 76.618 | 8.7402 | 1.572e-05 *** |
| period | 2 | 172.31 | 86.154 | 9.8279 | 0.0013030 ** |
| sequence | 5 | 318.69 | 63.738 | 7.2709 | 0.0006954 *** |
| trt | 2 | 440.61 | 220.304 | 25.1311 | 6.164e-06 *** |
| carry | 2 | 16.43 | 8.215 | 0.9372 | 0.4100385 |
| sequence:steer | 6 | 118.50 | 19.750 | 2.2530 | 0.0849122 . |
| RESIDUALS | 18 | 157.79 | 8.766 | | |
| CORRECTED TOTAL | 35 | 1460.31 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(y ~ period + sequence + steer:sequence + trt + carry, v1p410), type=3,
      singular.ok=TRUE) # NOT OK for sequence
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: y

| | Sum Sq | Df | F values | Pr(>F) |
|----------------|--------|----|----------|---------------|
| period | 172.31 | 2 | 9.8279 | 0.001303 ** |
| sequence | 0.00 | 0 | | |
| trt | 440.61 | 2 | 25.1311 | 6.164e-06 *** |
| carry | 16.43 | 2 | 0.9372 | 0.410038 |
| sequence:steer | 118.50 | 6 | 2.2530 | 0.084912 . |
| Residuals | 157.79 | 18 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8 Searle - Linear Models 2e

Reference

- Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

8.1 7.2 (p390, 59%)

(19) MODEL

```
weight = c(8,13,9,12,7,11,6,12,12,14,9,7,14,16,10,14,11,13)
treatment = c("ta","ta","ta","ta","ta","tb","tb","tb","tb","tc","tc","tc",
             "tc","tc","tc")
variety = c("va","va","va","vc","vd","vd","va","va","vb","vb","vb","vb",
           "vc","vd","vd","vd")
d1 = data.frame(weight, treatment, variety)
aov3(weight ~ treatment*variety, d1)
```

```
Response : weight
            Df  Sum Sq Mean Sq F value Pr(>F)
MODEL          7  82.000 11.7143  2.0918 0.13995
treatment      2   12.471  6.2353  1.1134 0.36595
variety        3   34.872 11.6240  2.0757 0.16719
treatment:variety  2   34.714 17.3571  3.0995 0.08965 .
RESIDUALS      10   56.000  5.6000
CORRECTED TOTAL 17  138.000
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(weight ~ treatment*variety, d1), type=3, singular.ok=TRUE) # NOT OK
```

```
Note: model has aliased coefficients
      sums of squares computed by model comparison
```

```
Anova Table (Type III tests)
```

```
Response: weight
            Sum Sq Df F values Pr(>F)
treatment      0.000  0
variety        0.000  0
treatment:variety 34.714  2   3.0995 0.08965 .
Residuals     56.000 10
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.2 7.2 (p393, 60%)

(20) MODEL

```
percent = c(31,33,44,36,38,26,37,59,42,42,34,42,28,39,36,32,38,42,36,22,42,46,
          26,37,43)
refinery = c(rep("g",9),rep("n",8),rep("s",8))
process = as.factor(c(1,1,1,1,1,1,2,2,2,1,1,1,2,2,2,2,1,1,1,2,2,2,2,2))
source0 = c("t","t","t","t","o","m","t","t","o","m","i","i","t","o","m","m",
           "t","o","i","o","o","m","i","i")
d2 = data.frame(percent, refinery, process, source=source0)
aov3(percent ~ refinery*source, d2)
```

Response : percent

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-----------------|----|---------|---------|---------|--------|
| MODEL | 10 | 442.56 | 44.256 | 0.6361 | 0.7616 |
| refinery | 2 | 10.77 | 5.383 | 0.0774 | 0.9259 |
| source | 3 | 282.63 | 94.211 | 1.3542 | 0.2972 |
| refinery:source | 5 | 155.47 | 31.095 | 0.4469 | 0.8086 |
| RESIDUALS | 14 | 974.00 | 69.571 | | |
| CORRECTED TOTAL | 24 | 1416.56 | | | |

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(percent ~ refinery*source, d2), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: percent

| | Sum Sq | Df | F values | Pr(>F) |
|-----------------|--------|----|----------|--------|
| refinery | 2.52 | 1 | 0.0362 | 0.8518 |
| source | 268.19 | 2 | 1.9275 | 0.1822 |
| refinery:source | 155.47 | 5 | 0.4469 | 0.8086 |
| Residuals | 974.00 | 14 | | |

9 Web site examples

9.1 <https://github.com/djnavarro/psyr>

(21) MODEL

```
d21 = read.csv("http://r.acr.kr/psyr/coffee.csv")
GLM(babble ~ sugar*milk - 1, d21)
```

\$ANOVA

Response : babble

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|-------------------|----|--------|---------|---------|--------------|
| MODEL | 6 | 472.54 | 78.756 | 298.84 | 2.39e-12 *** |
| RESIDUALS | 12 | 3.16 | 0.264 | | |
| UNCORRECTED TOTAL | 18 | 475.70 | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

| Root MSE | babble | Mean Coef | Var | R-square | Adj R-sq |
|-----------|----------|-----------|-----------|-----------|----------|
| 0.5133631 | 5.066667 | 10.13217 | 0.9933519 | 0.9900279 | |

\$`Type I`

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|------------|----|--------|---------|----------|---------------|
| sugar | 3 | 465.64 | 155.213 | 588.9486 | 2.756e-13 *** |
| milk | 1 | 0.96 | 0.956 | 3.6279 | 0.081061 . |
| sugar:milk | 2 | 5.94 | 2.972 | 11.2769 | 0.001754 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|------------|----|--------|---------|---------|-------------|
| sugar | 2 | 3.0696 | 1.53482 | 5.8238 | 0.017075 * |
| milk | 1 | 0.9561 | 0.95611 | 3.6279 | 0.081061 . |
| sugar:milk | 2 | 5.9439 | 2.97193 | 11.2769 | 0.001754 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

CAUTION: Singularity Exists !

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|------------|----|--------|---------|---------|-------------|
| sugar | 2 | 2.1318 | 1.0659 | 4.0446 | 0.045426 * |
| milk | 1 | 1.0041 | 1.0041 | 3.8102 | 0.074672 . |
| sugar:milk | 2 | 5.9439 | 2.9719 | 11.2769 | 0.001754 ** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
```

```
r21 = lm(babble ~ sugar*milk - 1, d21)
```

```
Anova(r21, type=2) # NOT OK
```

Anova Table (Type II tests)

Response: babble

| | Sum Sq | Df | F value | Pr(>F) |
|------------|--------|----|----------|---------------|
| sugar | 453.76 | 3 | 573.9233 | 3.214e-13 *** |
| milk | 0.96 | 1 | 3.6279 | 0.081061 . |
| sugar:milk | 5.94 | 2 | 11.2769 | 0.001754 ** |
| Residuals | 3.16 | 12 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
Anova(r21, type=3) # NOT OK
```

Anova Table (Type III tests)

Response: babble

| | Sum Sq | Df | F value | Pr(>F) |
|------------|--------|----|----------|---------------|
| sugar | 454.77 | 3 | 575.1970 | 3.172e-13 *** |
| milk | 1.00 | 1 | 3.8102 | 0.074672 . |
| sugar:milk | 5.94 | 2 | 11.2769 | 0.001754 ** |
| Residuals | 3.16 | 12 | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10 Bioequivalence (BE) data example

(22) MODEL

```
GLM(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata) # a BE dataset in sasLM package
```

\$ANOVA

```
Response : log(CMAX)
            Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL          48 23.1924 0.48317  5.6278 4.395e-08 ***
RESIDUALS      42  3.6059 0.08585
CORRECTED TOTAL 90 26.7983
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
Root MSE log(CMAX) Mean Coef Var R-square Adj R-sq
0.2930098      6.071036 4.826355 0.8654428 0.7116631
```

\$`Type I`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6454 0.64544  7.5178  0.008938 **
SEQ:SUBJ 45 22.4395 0.49866  5.8081 3.359e-08 ***
PRD     1  0.0969 0.09686  1.1281  0.294242
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6440 0.64395  7.5005  0.009011 **
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.287632
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.3368 0.33679  3.9228  0.05421 .
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.28763
TRT     1  0.0106 0.01057  0.1231  0.72741
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts=c("contr.sum", "contr.poly"))
```

```
Anova(lm(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata), type=3, singular.ok=TRUE)
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: log(CMAX)

| | Sum Sq | Df | F values | Pr(>F) |
|---|---------|----|----------|---------------|
| SEQ | 0.0000 | 0 | | |
| PRD | 0.0996 | 1 | 1.1599 | 0.2876 |
| TRT | 0.0106 | 1 | 0.1231 | 0.7274 |
| SEQ:SUBJ | 22.5232 | 45 | 5.8298 | 3.173e-08 *** |
| Residuals | 3.6059 | 42 | | |
| --- | | | | |
| Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1 | | | | |

11 Session Information

```
R version 4.3.1 (2023-06-16 ucrt)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 10 x64 (build 19044)
```

```
Matrix products: default
```

```
locale:
```

```
[1] LC_COLLATE=Korean_Korea.utf8  LC_CTYPE=Korean_Korea.utf8
[3] LC_MONETARY=Korean_Korea.utf8 LC_NUMERIC=C
[5] LC_TIME=Korean_Korea.utf8
```

```
time zone: Asia/Seoul
```

```
tzcode source: internal
```

```
attached base packages:
```

```
[1] stats      graphics   grDevices utils      datasets  methods   base
```

```
other attached packages:
```

```
[1] car_3.1-2       carData_3.0-5  sasLM_0.9.12  mvtnorm_1.2-3 rmarkdown_2.24
```

```
loaded via a namespace (and not attached):
```

```
[1] digest_0.6.33   fastmap_1.1.1   xfun_0.40      abind_1.4-5
[5] knitr_1.43     htmltools_0.5.6  cli_3.6.1      compiler_4.3.1
[9] tools_4.3.1    evaluate_0.21   yaml_2.3.7     rlang_1.1.1
[13] MASS_7.3-60
```