

# sstvars: Structural Smooth Transition Vector Autoregressive Models in R

Savi Virolainen  
University of Helsinki

---

## Abstract

We describe the R package **sstvars**, which provides tools for estimating and analyzing the reduced form and structural smooth transition vector autoregressive (STVAR) models. This includes also threshold vector autoregressive models. The package implements various transition weight functions, conditional distributions, identification methods, and parameter restrictions. The model parameters are estimated with the method of (penalized) maximum likelihood (ML) by running multiple rounds of a two-phase or a three-step procedure estimation procedure. In the two-phase procedure, a genetic algorithm is used in the first phase to find starting values for a gradient based variable metric algorithm, which is ran in the second phase. In the three-step procedure, autoregressive and weight parameters are first estimated by (nonlinear) least squares, then the rest of the parameters are estimated by ML conditional on them with a genetic algorithm, and finally all the parameters are estimated by (penalized) ML by initializing the variable metric algorithm from the initial estimates obtained from the first two steps. Both penalized and "normal" ML estimation are supported. For evaluating the adequacy of the estimated models, **sstvars** utilizes residuals based diagnostics and provides functions for graphical diagnostics and for calculating formal diagnostic tests. **sstvars** also accommodates the computation of linear impulse response functions, nonlinear generalized impulse response functions, generalized forecast error variance decompositions, and historical decompositions, as well as conducting counterfactual analysis. Further functionality includes hypothesis testing, plotting the profile log-likelihood functions about the estimate, simulation from STVAR processes, and forecasting, to name a few. We illustrate the use of **sstvars** with a quarterly series consisting of two U.S. variables: the percentage change of real GDP and the percentage change of GDP implicit price deflator.

*Keywords:* smooth transition vector autoregressive model, structural smooth transition vector autoregressive model, regime-switching, threshold VAR model, SVAR, TVAR, STVAR.

---

## Contents

<b>Introduction</b>	<b>1</b>
<b>Smooth Transition Vector Autoregressive Models</b>	<b>2</b>
General framework for STVAR models . . . . .	3
Stationarity condition . . . . .	4
Specifications of the conditional distribution . . . . .	5
Gaussian distribution . . . . .	5
Student's t distribution . . . . .	5
Independent Student's t distribution . . . . .	5
Independent skewed t distribution . . . . .	6
Specifications of the transition weights . . . . .	7
Weighted relative likelihoods . . . . .	7
Logistic transition weights . . . . .	7
Multinomial logit transition weights . . . . .	8
Exponential transition weights . . . . .	9
Threshold transition weights . . . . .	9
Exogenous transition weights . . . . .	10
<b>Structural STVAR models</b>	<b>10</b>
Recursive identification . . . . .	10
Identification via heteroskedasticity . . . . .	11
Identification by non-Gaussianity . . . . .	12
Identification of the shocks in TVAR models . . . . .	12
Identification of the shocks in STVAR models . . . . .	14
Labelling the shocks and blended identification . . . . .	14
<b>Estimation</b>	<b>15</b>
The log-likelihood function . . . . .	16
The penalized log-likelihood function . . . . .	17
Two-phase estimation procedure . . . . .	17
Three-step estimation procedure . . . . .	18
Examples of unconstrained estimation . . . . .	19
Further examination of the estimates . . . . .	24
Estimation of structural STVAR models . . . . .	27
Linear constraints on the autoregressive parameters . . . . .	29
Constraining the unconditional means of some regimes to be equal . . . . .	32
Constraining weight functions parameters . . . . .	32
<b>Testing parameter constraints</b>	<b>33</b>
<b>Residual based model diagnostics</b>	<b>33</b>

<b>Impulse response analysis</b>	<b>34</b>
Generalized impulse response function . . . . .	34
Generalized forecast error variance decomposition . . . . .	35
Linear impulse response functions . . . . .	36
<b>Historical decompositions</b>	<b>36</b>
<b>Counterfactual analysis</b>	<b>37</b>
Historical counterfactuals . . . . .	39
Forecast scenarios . . . . .	39
Counterfactual GIRFs . . . . .	41
<b>Building a STVAR model with specific parameter values</b>	<b>41</b>
<b>Simulation and forecasting</b>	<b>43</b>
Simulation . . . . .	43
Simulation based forecasting . . . . .	43
<b>Summary</b>	<b>44</b>
<b>A table of some useful functions</b>	<b>45</b>

## 1. Introduction

Linear vector autoregressive (VAR) models are a standard tools in time series econometrics. They can be employed to answer questions about the statistical relationships of different variables or to forecast future values of the process, for example. Structural VAR models, in particular, allow to trace out the effects of economic shocks. With an appropriate choice of the autoregressive order  $p$ , a linear VAR model is often able to filter out autocorrelation from the series very well. If the errors are assumed to follow an autoregressive conditional heteroskedasticity (ARCH) process, the model is also often able to adequately filter out conditional heteroskedasticity from the series.

In some cases, linear VAR models are not, however, capable of capturing all the relevant characteristics of the data. This includes shifts in the mean or volatility, and changes in the autoregressive dynamics of the process. Such nonlinear features frequently occur in economic time series when the underlying data generating dynamics vary in time, for example, depending the specific state of the economy. Various types of time series models capable of capturing this kind of regime-switching behavior have been proposed, one of them is the smooth transition vector autoregressive (STVAR) models that allow to capture gradual shifts in the dynamics of the data. They consist of a finite number of regimes, each of which are linear vector autoregressions that are characterized by different autoregressive coefficients or error term covariance matrices. The package **sstvars** considers STVAR models in which, at each point of time, the observation is a weighted average of the conditional means of the regimes plus a random error whose covariance matrix is a weighted average of the covariance matrices of the regimes. The weights, in turn, are expressed in terms of time-varying transition weights that either depend on the preceding observations or are exogenous. Different STVAR models can be created by specifying the transition weights or the error distribution in various ways.

This manuscript describes the R package **sstvars** providing a set of easy-to-use tools for STVAR modeling, including also threshold vector autoregressive models. There are tools for unconstrained and constrained (penalized) maximum likelihood (ML) estimation of the model parameters, residual based model diagnostics, counterfactual analysis, as well as for computation of linear impulse response functions, nonlinear generalized impulse response functions, generalized forecast error variance decompositions, and historical decompositions. In addition to "normal" ML estimation, **sstvars** also accommodates penalized ML estimation, in which a penalization term is added to the log-likelihood function, penalizing parameter values that are inside or are close entering instability region (where the usual stability condition does not hold). This enables to allow for unstable parameter values in estimation, letting the optimization algorithm to explore the unstable region of the parameter space, often substantially improving its performance. By penalizing instability sufficiently, the optimization algorithm eventually steers back to the stable region. Further functionality includes hypothesis testing, plotting the profile log-likelihood functions about the estimate, simulation from STVAR processes, and forecasting, to name a few. Various transition weight functions are accommodated, including exogenous weights, logistic weights (Anderson and Vahid 1998), multinomial logit weights, exponential weights (e.g., Hubrich and Teräsvirta 2013), threshold weights (Tsay 1998), and transition weights that defined as weighted relative likelihoods of the regimes corresponding to the preceding  $p$  observations (Lanne and Virolainen 2025). Currently, the accommodated conditional distributions include Gaussian distribution, Student's  $t$  distribution, Student's  $t$  distribution with mutually independent components, and skewed  $t$  distribution with mutually independent components. The accommodated identification methods include recursive identification, identification by heteroskedasticity (Lütkepohl and Netšunajev 2017), and identification by non-Gaussianity (Virolainen 2025a).

The estimation of the model parameters can, in some cases, be rather tricky. Particularly when the transition weights are determined endogenously, there is a very large number of modes

to the (penalized) log-likelihood function, and large areas of the parameter space, where the (penalized) log-likelihood function is flat in multiple directions. Therefore, the model parameters are estimated by running multiple rounds of a two-phase or three phase estimation procedure. In the two-phase procedure, a genetic algorithm is used in the first phase to find starting values for a gradient based variable metric algorithm, which is ran in the second phase. In the three-step procedure, autoregressive and weight parameters are first estimated by (nonlinear) least squares (LS), then the rest of the parameters are estimated by (penalized) ML conditional on them with a genetic algorithm, and finally all the parameters are estimated by (penalized) ML by initializing the variable metric algorithm from the initial estimates obtained from the first two steps. Because of the multimodality of the (penalized) log-likelihood function, some of the estimation rounds may end up in different local maximum points, thereby enabling the researcher to build models not only based on the global maximum point but also on the local ones. The estimated models can be conveniently examined with the `summary` and `plot` methods.

The three-step procedure facilitates estimating larger model and substantially reduces to computation time as opposed to the two-phase procedure, as a substantially smaller number of estimation rounds are required due to the initial estimates from (nonlinear) LS estimation. However, the three-step procedure tends to produce estimates that are close to the initial (N)LS estimates, while the two-phase procedure explores the parameter space more thoroughly (when a large enough number of estimation rounds is ran). The penalized ML estimation is particularly useful when the data is persistent (which macroeconomic data typically is), as it allows the optimization algorithm to explore the unstable region of the parameter space (including the boundary), often improving its performance substantially. Moreover, penalized estimation can more effectively make use of (N)LS estimates that are outside the stability region, as with penalized estimation they do not need to be adjusted when switching to ML estimation in the latter phases.

The remainder of this paper is organized as follows. Section 2 defines the implemented reduced form STVAR models and discusses some of their properties. Section 3 discusses the accommodated structural STVAR models and discusses identification of the structural shocks. Section 4 discusses estimation of the model parameters. We also illustrate how the STVAR models can be estimated and examined with `sstvars` and how various parameter restrictions can be imposed in the estimation. Section 6 discusses how to evaluate the model adequacy with `sstvars` using residual based diagnostics. Section 7 discusses impulse response analysis, including generalized impulse response functions and generalized forecast error variance decompositions. Section 8 discusses computation of historical decompositions and Section 9 counterfactual analysis. Section 10 shows how the STVAR models can be constructed with given parameter values. In Section 11, we first show how to simulate observations from a STVAR process, and then we illustrate how to forecast future values of a STVAR process with a simulation-based Monte Carlo method. Finally, Section 12 concludes, and some useful functions in `sstvars` are collected to a single table in Section 12 for convenience. Throughout this paper, we illustrate the use of `sstvars` with a quarterly series consisting of two U.S. variables: the percentage change of real GDP and the percentage change of GDP implicit price deflator, covering the period from 1959Q1 to 2019Q4.

## 2. Smooth Transition Vector Autoregressive Models

This section describes the STVAR models implemented in `sstvars`. First, we describe the general framework of STVAR models accommodated by `sstvars` and present a sufficient condition for their ergodic stationarity. Then, we present the implemented specifications of transition weight functions and conditional distributions. Finally, we discuss structural STVAR models and implemented methods for identification of the shocks.

## 2.1. General framework for STVAR models

Let  $y_t$ ,  $t = 1, 2, \dots$ , be the  $d$ -dimensional time series of interest and  $\mathcal{F}_{t-1}$  denote the  $\sigma$ -algebra generated by the random vectors  $\{y_{t-j}, j > 0\}$ . We consider STVAR models with  $M$  regimes and autoregressive order  $p$  assumed to satisfy

$$y_t = \sum_{m=1}^M \alpha_{m,t} \mu_{m,t} + u_t, \quad u_t \sim MD(0, \Omega_{y,t}) \quad (1)$$

$$\mu_{m,t} = \phi_m + \sum_{i=1}^p A_{m,i} y_{t-i}, \quad m = 1, \dots, M, \quad (2)$$

where  $\phi_m \in \mathbb{R}^d$  are intercept parameters,  $A_{m,i}$  is the lag  $i$  autoregression matrix of Regime  $m$ , and  $u_t$  is a martingale difference sequence of reduced form innovations. The transition weights  $\alpha_{m,t}$  are assumed to be  $\mathcal{F}_{t-1}$ -measurable functions of  $\{y_{t-j}, j = 1, \dots, p\}$  and to satisfy  $\sum_{m=1}^M \alpha_{m,t} = 1$  at all  $t$ . They express the proportions of the regimes the process is on at each point of time, and how the process shifts between the regimes. Through, a STVAR model with autoregressive order  $p$  and  $M$  regimes is referred to as a STVAR( $p, M$ ) model, whenever the order of the model needs to be emphasized.

Conditional on  $\mathcal{F}_{t-1}$ , the conditional mean of the above described process is  $\mu_{y,t} \equiv E[y_t | \mathcal{F}_{t-1}] = \sum_{m=1}^M \alpha_{m,t} \mu_{m,t}$ . The conditional mean is thereby a weighted sum the regime-specific means  $\mu_{m,t}$  with the weights given by the transition weights  $\alpha_{m,t}$ . The specification of the conditional covariance matrix  $\Omega_{y,t} \equiv \text{Cov}(y_t | \mathcal{F}_{t-1}) = \text{Cov}(u_t | \mathcal{F}_{t-1})$  depends on the error term distribution. If the conditional distribution of  $u_t$  is either Gaussian or Student's  $t$ -distribution, the conditional covariance matrix is assumed to be the weighted average of the covariance matrices of the regimes:

$$\Omega_{y,t} = \sum_{m=1}^M \alpha_{m,t} \Omega_m, \quad (3)$$

where  $\Omega_1, \dots, \Omega_M$  are the positive definite ( $d \times d$ ) covariance matrices of the regimes.

If the distribution of  $u_t$  is the Student's  $t$ -distribution with mutually independent components (hereafter independent  $t$ -distribution) or independent skewed  $t$ -distribution with mutually independent components (see Hansen 1994), the conditional covariance matrix is different, because in that case the model is directly parametrized with the invertible ( $d \times d$ ) impact matrices  $B_m$  of the regimes as:

$$y_t = \sum_{m=1}^M \alpha_{m,t} \mu_{m,t} + B_{y,t} e_t, \quad e_t \sim IID(0, I_d) \quad (4)$$

$$B_{y,t} = \sum_{m=1}^M \alpha_{m,t} B_m, \quad (5)$$

where  $B_1, \dots, B_M$  are the invertible ( $d \times d$ ) impact matrices of the regimes and  $e_t$  is the sequence of structural errors. See Virolainen (2025a) for further details.

Different STVAR models are obtained by specifying the transition weights or the error distribution (i.e., the conditional distribution) in various ways. See Hubrich and Teräsvirta (2013) for a survey on STVAR literature, including formulations more general than our framework. The package **sstvars** accommodates completely exogenous transition weight functions as well as transition weights that are functions of  $\{y_{t-j}, j = 1, \dots, p\}$ . In the latter case, the stationarity condition presented below applies. Moreover, if the transition weights are not exogenous, their  $\mathcal{F}_{t-1}$ -measurability ensures that the true generalized impulse responses functions (Koop, Pesaran, and Potter 1996) can be easily estimated, as completely exogenous switching-variables are excluded from affecting the weights. We also assume that the transition weights are identical

for all the individual equations in (1), which is also required for applicability of the stationarity condition. Consequently, at each  $t$ , the process can be described as a weighted sum of linear vector autoregressions.

## 2.2. Stationarity condition

Excluding models with exogenous transition weights, it can be shown that a sufficient condition for the ergodic stationarity of the STVAR model (1)-(2) can be expressed in terms of the joint spectral radius (JSR) of certain matrices (Kheifets and Saikkonen 2020). The JSR of a finite set of square matrices  $\mathcal{A}$  is defined by

$$\rho(\mathcal{A}) = \limsup_{j \rightarrow \infty} \left( \sup_{A \in \mathcal{A}^j} \rho(A) \right)^{1/j}, \quad (6)$$

where  $\mathcal{A}^j = \{A_1 A_2 \dots A_j : A_i \in \mathcal{A}\}$  and  $\rho(A)$  is the spectral radius of the square matrix  $A$ .

Consider the companion form AR matrices of the regimes defined as

$$\mathbf{A}_m = \begin{bmatrix} A_{m,1} & A_{m,2} & \cdots & A_{m,p-1} & A_{m,p} \\ I_d & 0 & \cdots & 0 & 0 \\ 0 & I_d & & 0 & 0 \\ \vdots & & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & I_d & 0 \end{bmatrix}, \quad m = 1, \dots, M. \quad (7)$$

$(dp \times dp)$

Kheifets and Saikkonen (2020, Theorem 1) and Lanne and Virolainen (2025) (see also Saikkonen 2008) show that if the following condition holds, the STVAR process is ergodic stationary (both strictly and second-order).

**Condition 1**  $\rho(\{\mathbf{A}_1, \dots, \mathbf{A}_M\}) < 1$ .

Condition 1 is, however, computationally demanding the check in practice with a reasonable accuracy (e.g., Chang and Blondel 2013), making it impractical to use in the estimation. Therefore, we consider a necessary condition for Condition 1 that is easier to check in practice, which is that the usual stability condition is satisfied for each of the regimes. Specifically, the following condition, which is analogous to Corollary 1 of Kheifets and Saikkonen (2020), is necessary for Condition 1.

**Condition 2**  $\max\{\rho(\mathbf{A}_1), \dots, \rho(\mathbf{A}_M)\} < 1$ ,

where  $\rho(\mathbf{A}_m)$  is the spectral radius of  $\mathbf{A}_m$ ,  $m = 1, \dots, M$ .

Note that validity Condition 2 does not imply the validity of Condition 1, which guarantees ergodic stationarity of the model. However, in practice models that satisfy Condition 2 and are not very close to breaking this condition usually satisfy Condition 1. For checking the validity of Condition 1, `sstvars` implements (the function `bound_JSR`) implements the branch-and-bound method by Gripenberg (1996). For large models `sstvars`'s implementation of the Gripenberg's method may, however, take very long if tight bounds are required. Other implementations of methods bounding the JSR include the MATLAB toolbox `JSR` by Jungers (2023), which automatically combines several methods and finds accurate bounds much faster than `sstvars`.

The R package `sstvars` operates largely by assuming that all of the regimes satisfy the usual stability condition. However, also estimates that do not satisfy the usual stability condition are allowed in `sstvars`. In this case, the argument `allow_unstab` should be set to `TRUE`.

### 2.3. Specifications of the conditional distribution

Currently, **sstvars** accommodates two types of error distributions: Gaussian distribution, Student's  $t$ , and independent Student's  $t$  distribution, which are discussed below.

#### *Gaussian distribution*

Assuming the structural errors  $e_t$  have standard normal distributions, the conditional distribution of  $y_t$  conditional on  $\mathcal{F}_{t-1}$  is Gaussian and characterized by the density function

$$f(y_t|\mathcal{F}_{t-1}) = n_d(y_t; \mu_t, \Omega_t) = (2\pi)^{-d/2} \det(\Omega_t)^{-1/2} \exp \left\{ -\frac{1}{2} (y_t - \mu_t)' \Omega_t^{-1} (y_t - \mu_t) \right\}. \quad (8)$$

That is, the conditional distribution is simply the  $d$ -dimensional Gaussian distribution with mean  $\mu_t$  and covariance matrix  $\Omega_t$ . The Gaussian distribution simple and can be used with all of our transition weight functions, but in some cases it is useful to employ the more heavy tailed Student's  $t$  distribution instead.

#### *Student's $t$ distribution*

To accommodate more heavy tailed data, instead of using Gaussian errors one may consider Student's  $t$  errors and assume the shocks  $e_t$  are Student's  $t$  distributed with the mean zero, identity covariance matrix, and  $\nu > 2$  degrees of freedom (where the assumption  $\nu > 2$  is made to ensure the existence of second moments). The Student's  $t$  STVAR model has the conditional distribution, conditional  $\mathcal{F}_{t-1}$ , characterized by the density function

$$f(y_t|\mathcal{F}_{t-1}) = t_d(y_t; \mu_t, \Omega_t, \nu) = C_d(\nu) \det(\Omega_t)^{-1/2} \left( 1 + \frac{(y_t - \mu_t)' \Omega_t^{-1} (y_t - \mu_t)}{\nu - 2} \right)^{-(d+\nu)/2}, \quad (9)$$

where

$$C_d(\nu) = \frac{\Gamma\left(\frac{d+\nu}{2}\right)}{\sqrt{\pi^d (\nu - 2)^d} \Gamma\left(\frac{\nu}{2}\right)}, \quad (10)$$

and  $\Gamma(\cdot)$  is the gamma function. The conditional distribution is, hence, the  $d$ -dimensional Student's  $t$  distribution with mean  $\mu_t$ , covariance matrix  $\Omega_t$ , and  $\nu$  degrees of freedom. Note that the parametrization differs from the conventional one, as the distribution is parametrized with a covariance matrix instead a scale matrix (see, e.g., [Meitz, Preve, and Saikkonen 2023](#), Appendix A for details about the parametrization).

The Student's  $t$  errors are more flexible than the Gaussian ones, but they cannot be used with the transition weight function that is defined as weighted ratios of the regime's stationary densities (see Section 2.4.1). This is because it requires the knowledge of the stationary distributions of the regimes corresponding to  $p$  consecutive observations, and the stationary distribution is not known for the Student's  $t$  regimes. Moreover, Student's  $t$  STVAR models are more difficult estimate in practice than the Gaussian ones, and estimation of the STVAR models can be demanding.

#### *Independent Student's $t$ distribution*

In addition to the conventional multivariate Student's  $t$  distribution, **sstvars** accommodates Student's  $t$  distribution with mutually independent components, i.e., each component of the error term follows a univariate Student's  $t$  distribution independently from the the other components. The independent Student's  $t$  STVAR model has the conditional distribution, conditional  $\mathcal{F}_{t-1}$ , characterized by the density function

$$f(y_t|\mathcal{F}_{t-1}) = |\det(B_t)|^{-1} \prod_{i=1}^d t_1(\iota_i' B_t^{-1} (y_t - \mu_t); 0, 1, \nu_i) \quad (11)$$



where  $\iota_i = (0, \dots, 0, 1, 0, \dots, 0)$  is a  $(d \times 1)$  has one in the  $i$ th entry and zeros in the other entries,  $t_1(\cdot; 0, 1, \nu_i)$  is the density function of univariate  $t$ -distribution with mean zero, variance one, and  $\nu_i$  degrees of freedom (obtained from the  $d$ -dimensional density described in Section 2.3.2), and

$$B_{y,t} = \sum_{m=1}^M \alpha_{m,t}^{1/2} B_m, \quad (12)$$

where  $B_1, \dots, B_M$  are invertible  $(d \times d)$  impact matrices of the regimes.

The independent Student's  $t$  distribution is more flexible than the conventional  $t$ -distribution, as it allows for different degrees of freedom parameter values for each component. However, the for structural analysis, a more substantial advantage is that under mutually independent Student's  $t$  shocks, the structural shocks are statistically identified without further restrictions on the model (Virolainen 2025a). But this comes at a cost: due to the structure of the conditional distribution, evaluation of the log-likelihood function is computationally more costly than with the conventional  $t$ -distribution. Therefore, maximum likelihood estimation of STVAR with independent Student's  $t$  shocks is somewhat slower.

### Independent skewed $t$ distribution

We also accommodate the skewed  $t$  distribution with mutually independent components as conditional distribution. In other words, each shock  $e_{it}$  in the structural error  $e_t = (e_{1t}, \dots, e_{dt})$  is assumed to follow a univariate skewed  $t$  distribution described in Hansen (1994, Section 2.4) independently from the other shocks. The independent skewed  $t$  STVAR model has the conditional distribution, conditional  $\mathcal{F}_{t-1}$ , characterized by the density function

$$f(y_t | \mathcal{F}_{t-1}) = |\det(B_t)|^{-1} \prod_{i=1}^d st(\iota_i' B_t^{-1}(y_t - \mu_t); \nu_i, \lambda_i), \quad (13)$$

where  $\iota_i = (0, \dots, 0, 1, 0, \dots, 0)$  is a  $(d \times 1)$  has one in the  $i$ th entry and zeros in the other entries,  $B_{y,t}$  is as in (12), and  $st(\cdot; \nu_i, \lambda_i)$  is the density function of univariate skewed  $t$ -distribution with zero mean, unit variance,  $\nu_i$  degrees of freedom, and skewness controlled by the parameter  $\lambda_i \in (-1, 1)$ . The density function of the skewed  $t$ -distribution,  $st(\cdot; \nu_i, \lambda_i)$ , is given as (Hansen 1994, Equations (10)-(13)):

$$st(e_{it}; \nu_i, \lambda_i) = b_i c_i \left( 1 + \frac{1}{\nu_i - 2} \left( \frac{b_i e_{it} + a_i}{1 - \mathbf{1}\{e_{it} < -a_i/b_i\} \lambda_i + \mathbf{1}\{e_{it} \geq -a_i/b_i\} \lambda_i} \right)^2 \right)^{-(\nu_i+1)/2} \quad (14)$$

where  $\mathbf{1}\{e_{it} < -a_i/b_i\}$  is an indicator function that takes the value one if  $e_{it} < -a_i/b_i$  and zero otherwise, and  $\mathbf{1}\{e_{it} \geq -a_i/b_i\}$  is an indicator function that takes the value one if  $e_{it} \geq -a_i/b_i$  and zero otherwise. The constants  $a_i, b_i$ , and  $c_i$  are defined as

$$a_i = 4\lambda_i c_i \left( \frac{\nu_i - 2}{\nu_i - 1} \right), \quad (15)$$

$$b_i = (1 + 3\lambda_i^2 - a_i^2)^{1/2}, \quad \text{and} \quad (16)$$

$$c_i = \frac{\Gamma\left(\frac{\nu_i+1}{2}\right)}{(\pi(\nu_i - 2))^{1/2} \Gamma\left(\frac{\nu_i}{2}\right)}, \quad (17)$$

where  $\Gamma(\cdot)$  is Gamma function.

Independent skewed  $t$  distribution accommodates the independent Student's  $t$  distribution as the special case  $\lambda_i = 0$ . The independent skewed  $t$  distribution is, however, more flexible than the independent Student's  $t$  distribution, as it allows for skewed distributions for each component. On the other hand, models incorporating skewed shocks can be substantially more

difficult to estimate in practice, because of the increased complexity and since the estimation algorithm might have difficulties in distinguishing between skewness and shifts in the mean.

## 2.4. Specifications of the transition weights

Various specifications of the transition weights  $\alpha_{m,t}$  can be considered to obtain smooth transition VARs with different properties. We assume that the transition weights are either completely exogenous or functions of  $\{y_{t-j}, j = 1, \dots, p\}$ . Under the latter type of weights, the stationarity condition discussed in Section 2.2 applies. Moreover,  $\mathcal{F}_{t-1}$ -measurability of the transition weights ensures that the true generalized impulse responses functions can be easily estimated, as completely exogenous switching-variables are excluded from affecting the weights.<sup>1</sup> We also assume that the transition weights are identical for all the individual equations in (1), which is required for applicability of the stationarity condition. Consequently, at each  $t$ , the process can be described as a weighted sum of linear VARs.

### *Weighted relative likelihoods*

If the conditional distribution is specified to be Gaussian, weighted relative likelihoods of the regimes can be used as transition weights (Lanne and Virolainen 2025). In this specification, the transitions weights depend on the full distribution of the preceding  $p$  observations, they are defined identically to the mixing weights in the Gaussian mixture vector autoregressive (GMVAR) model of Kalliovirta, Meitz, and Saikkonen (2016). Denoting  $\mathbf{y}_{t-1} = (y_{t-1}, \dots, y_{t-p})$ , the transition weights are defined as

$$\alpha_{m,t} = \frac{\alpha_m n_{dp}(\mathbf{y}_{t-1}; \mathbf{1}_p \otimes \mu_m, \Sigma_{m,p})}{\sum_{n=1}^M \alpha_n n_{dp}(\mathbf{y}_{t-1}; \mathbf{1}_p \otimes \mu_n, \Sigma_{n,p})}, \quad m = 1, \dots, M, \quad (18)$$

where  $\alpha_1, \dots, \alpha_M$  are transition weight parameters that satisfy  $\sum_{m=1}^M \alpha_m = 1$  and  $n_{dp}(\cdot; \mathbf{1}_p \otimes \mu_m, \Sigma_{m,p})$  is the density function of the  $dp$ -dimensional Gaussian distribution with mean  $\mathbf{1}_p \otimes \mu_m$  and covariance matrix  $\Sigma_{m,p}$ . The symbol  $\mathbf{1}_p$  denotes a  $p$ -dimensional vector of ones,  $\otimes$  is Kronecker product,  $\mu_m = (I_d - \sum_{i=1}^p A_{m,i})^{-1} \phi_m$ , and the covariance matrix  $\Sigma_{m,p}$  is given in Lütkepohl (2005, Equation (2.1.39)), but using the parameters of the  $m$ th regime. That is,  $n_{dp}(\cdot; \mathbf{1}_p \otimes \mu_m, \Sigma_{m,p})$  corresponds to the density function of the stationary distribution of the  $m$ th regime.

The transition weights are thus weighted ratios of the stationary densities of the regimes corresponding to the preceding  $p$  observations. This specification is appealing, as it states that the greater the weighted relative likelihood of a regime is, the greater the weight of this regime is. The regimes are, hence, formed based on the statistical properties of the data and are not affected by the choice of the switching variables similarly to the logistic weights.

In the GMVAR model (Kalliovirta *et al.* 2016), the definition of the mixing weights also leads to attractive theoretical properties such as the knowledge of the stationary distribution of  $p + 1$  consecutive observations. But this is not the case in our STVAR model, as the structure of the model is different. The GMVAR model has been implemented to the R package **gmvar** (Virolainen 2018a), which works quite similarly to **sstvars**.

### *Logistic transition weights*

A common specification assumes logistic transition weights (e.g., Anderson and Vahid 1998) that vary according to the level of the switching variable, which we assume to be a lagged endogenous variable. Here we assume that the model has only two regimes ( $M = 2$ ), and in

<sup>1</sup>Weaker forms of exogeneity of specific variables can also be imposed by constraining the AR matrices  $A_{m,i}$ ,  $m = 1, \dots, M$ ,  $i = 1, \dots, p$ , or the impact matrix  $B_t$  accordingly (see Section 4.8).

the next section, we show how the logistic weights generalize to multinomial logit weights that accommodate more regimes.

The logistic transition weights are defined as

$$\begin{aligned}\alpha_{1,t} &= 1 - \alpha_{2,t}, \\ \alpha_{2,t} &= [1 + \exp\{-\gamma(y_{it-j} - c)\}]^{-1},\end{aligned}\tag{19}$$

where  $y_{it-j}$  is the  $j$ th lagged observation ( $j \in \{1, \dots, p\}$ ) of the  $i$ th variable ( $i \in \{1, \dots, d\}$ ),  $c \in \mathbb{R}$  is a location parameter, and  $\gamma > 0$  is a scale parameter. The location parameter  $c$  determines the mid point of the transition function, i.e., the value of the (lagged) switching variable when the weights are equal. The scale parameter  $\gamma$ , in turn, determines the smoothness of the transitions (smaller  $\gamma$  implies smoother transitions), and it is assumed strictly positive so that  $\alpha_{2,t}$  is increasing in  $y_{it-j}$ .

Compared to weighted relative likelihoods, an advantage of the logistic weights is that it allows to specify switching variables in a way that leads to the regimes the econometrician is interested in in a specific application. For instance, if one is interested in how the effects of the shocks vary along with business cycle fluctuations,  $y_{it-j}$  may be set as a lagged output gap variable. STVAR models with logistics weights are also easier to estimate than those with the transition weights determined by weighted relative likelihoods of the regimes. A disadvantage is that the empirical results depend highly on the choice of the switching variable, and only the level of the switching variable affects the transition weights.

### *Multinomial logit transition weights*

The logistic transition weights can be generalized to multinomial logit weights that accommodate more than two regimes as well as multiple lags of multiple switching variables as regressors in the logit sub model. The generality, however, comes at the cost of significantly more difficult estimation in the practice and loss of the intuitive interpretations of the parameters of the transition function. With  $M \geq 2$  regimes, we specify the multinomial logit weights as

$$\alpha_{m,t} = \frac{\exp\{\gamma'_m z_{t-1}\}}{\sum_{n=1}^M \exp\{\gamma'_n z_{t-1}\}}, \quad m = 1, \dots, M,\tag{20}$$

where  $z_{t-1}$  is an  $(k \times 1)$   $\mathcal{F}_{t-1}$ -measurable vector containing the (lagged) switching variables and a constant term,  $\gamma_m$ ,  $m = 1, \dots, M - 1$ , are  $(k \times 1)$  coefficient vectors, and the last one is normalized as  $\gamma_M = 0$  ( $k \times 1$ ) to facilitate identification.<sup>2</sup> Denote the set of switching variables as  $I \subset \{1, \dots, d\}$  (with the indices in  $I$  corresponding to the ordering of the variables in  $y_t$ ) and assume that  $\tilde{p} \in \{1, \dots, p\}$  lags are included in the transition weights. We assume

$$z_{t-1} = (1, \tilde{z}_{\min\{I\}}, \dots, \tilde{z}_{\max\{I\}}), \quad \tilde{z}_j = (y_{it-1}, \dots, y_{it-\tilde{p}}), \quad i \in I.\tag{21}$$

So  $k = 1 + |I|\tilde{p}$  where  $|I|$  is the cardinality of the set  $I$  (i.e., the number of elements in  $I$ ). For instance, if the switching variables are the first two variables in  $y_t$ ,  $I = \{1, 2\}$  and only the first lag is included,  $\tilde{p} = 1$ , we have  $z_{t-1} = (1, y_{1t-1}, y_{2t-1})$ .

The specification implies

$$\log \frac{\alpha_{m,t}}{\alpha_{M,t}} = \gamma'_m z_{t-1}, \quad m = 1, \dots, M - 1.\tag{22}$$

Our specification assumes that all the lags up to the lag  $\tilde{p}$  are included in the transition weights. The inclusion of only some specific lags in the transition weights is, however, accommodated by

<sup>2</sup>Burgard, Neuenkirch, and Nöckel (2019) specify the mixing weights of their mixture VAR in a similar fashion, but unlike us, they allow for exogenous switching variables.

imposing constraints on the parameters  $\alpha \equiv (\gamma_1, \dots, \gamma_{M-1})$   $((M-1)k \times 1)$ . Specifically, **sstvars** assumes constraints of the form

$$\alpha = R\xi + r, \quad (23)$$

where  $R$  is a known  $((M-1)k \times l)$  constraint matrix,  $r$  is a known  $((M-1)k \times 1)$  constant, and  $\xi$  is an unknown  $(l \times 1)$  parameter. For instance, by assuming that  $R$  is a matrix of zeros, the weight parameter  $\alpha$  can be constrained to a known constant.

The logistic weights discussed in the previous section, with  $y_{it-j}$  as the switching variable for some lag  $j \in \{1, \dots, p\}$  and  $i \in \{1, \dots, d\}$ , are obtained as a special case as follows. Assume  $M = 2$ ,  $\tilde{p} = j$ , and  $I = \{i\}$ , so that  $z_{t-1} = (1, y_{it-1}, \dots, y_{it-j})$ . Then, impose the constraints  $r = 0$  and

$$R = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 1 \end{bmatrix} \quad (j+1 \times 2), \quad (24)$$

so  $\xi = (\gamma_{1,1}, \gamma_{j+1,1})$ , where  $\gamma_{l,m}$  is the  $l$ th element of  $\gamma_m$ .

A direct calculation shows that the "scale parameter" is  $-\gamma_{j+1,1}$  and the "location parameter" is  $\frac{\gamma_{1,1}}{-\gamma_{j+1,1}}$ . The linear constraints (23) do not, however, enable to constrain the location parameter  $\frac{\gamma_{1,1}}{-\gamma_{j+1,1}}$  to a specific value while leaving the scale parameter  $-\gamma_{(j+1),1}$  unconstrained (or vice versa), or to constrain the scale parameter strictly positive. Therefore, it is more convenient to use the logistic weights and parametrization discussed in Section 2.4.2 when only two regimes and one lag of one switching variable are used. Also, due to the structure of the weights, the multinomial logit weights are more difficult to estimate than the logistic weights.

### *Exponential transition weights*

Exponential transition weights (see, e.g., [Teräsvirta 1994](#)) vary according to the level of the switching variable, which we assume to be a lagged endogenous variable. Similarly to the logistic transition weights discussed in Section 2.4.2, the exponential weights depend on a location parameter  $c$  and a scale parameter  $\gamma$  that determine the mid point of the transition curve and smoothness of the transitions, respectively. But instead of logistic transition function, we consider an exponential transition function.

Specifically, we assume  $M = 2$  and define the exponential transition weights as

$$\begin{aligned} \alpha_{1,t} &= 1 - \alpha_{2,t}, \\ \alpha_{2,t} &= 1 - \exp\{-\gamma(y_{it-j} - c)^2\} \end{aligned} \quad (25)$$

where  $y_{it-j}$  is the  $j$ th lagged observation ( $j \in \{1, \dots, p\}$ ) of the  $i$ th variable ( $i \in \{1, \dots, d\}$ ),  $c \in \mathbb{R}$  is a location parameter, and  $\gamma > 0$  is a scale parameter. The location parameter  $c$  determines the value of the (lagged) switching variable when the process is completely in first regime, i.e.,  $\alpha_{1,t} = 1$  and  $\alpha_{2,t} = 0$ . The closer  $y_{it-j}$  is to  $c$ , the greater the weight of the first regime is. Conversely, when the deviation of  $y_{it-j}$  from  $c$  increases, the weight of the second regime increases (and the weight of the first regime decreases). The scale parameter  $\gamma$ , in turn, determines the smoothness of the transitions (smaller  $\gamma$  implies smoother transitions), and it is assumed strictly positive so that  $\alpha_{2,t} \in [0, 1]$  for all  $y_{it-j}$ .

### *Threshold transition weights*

Threshold transition weights assume discrete regime switches such that the regime switches when the level of the switching variable exceeds or falls below a threshold value. This type model nonlinear VARs are often referred to as Threshold VAR (TVAR) models ([Tsay 1998](#))

or self-exciting TVAR models (due to the endogenous switching-variable). We interpret the TVAR model as a special case of the STVAR models, despite of the regime switches being discrete rather than smooth.

For a model with  $M > 1$  regimes, consider the  $M-1$  threshold values  $r_1, \dots, r_{M-1} \in \mathbb{R}$  such that  $r_1 < \dots < r_{M-1}$ , and suppose the switching variable is  $i \in \{1, \dots, d\}$  with the lag  $j \in \{1, \dots, p\}$ . The transition function is defined as

$$\alpha_{m,t} = \begin{cases} 1 & \text{if } r_{m-1} < y_{it-j} \leq r_m, \\ 0 & \text{otherwise,} \end{cases} \quad (26)$$

where  $r_0 \equiv -\infty$ ,  $r_M \equiv \infty$ , and  $m = 1, \dots, M$ . In other words, at each  $t$ , the model defined in Equations (1), (2) and (26) reduces to a linear VAR corresponding to one of the regimes that is determined according to the level of the switching variable  $y_{it-j}$ .

Compared to smoothly varying transition weights, threshold transition weights have the advantage that the resulting model is easier to estimate in practice and the regimes have clearer interpretations. An obvious disadvantage is the inability to capture gradual shifts between the regimes or more complex regime-switching dynamics that depend on other factors than just on the level of the switching variable.

### *Exogenous transition weights*

In addition to the endogenous transition weights described above, **sstvars** accommodates completely exogenous transition weights. These weights are specified by the user by supplying them as a matrix. The only restrictions are that they must sum to one for each time period  $t$  and they must be weakly larger than zero. If exogenous weights are provided, the stationarity condition does not apply, but **sstvars** still assumes that each of the regimes satisfies the usual stability condition. Moreover, computation of the generalized impulse response functions requires that the exogenous transition weights that should be used for the sample paths are provided by the user. Linear impulse response functions based on a specific regime can, nonetheless, be calculated.

## 3. Structural STVAR models

Constructing a structural STVAR model from a reduced form STVAR model amounts to identifying the mutually and serially uncorrelated structural shocks  $e_t = (e_{t1}, \dots, e_{td})$ . The structural shocks are recovered from the reduced form innovations  $u_t$  based on the identity  $e_t = B_t^{-1}u_t$ , where  $B_t$  is a time-varying invertible  $(d \times d)$  impact matrix that governs the contemporaneous relationships of the shocks. Since many different solutions to the impact matrix generally lead to observationally equivalent models, further assumptions are required for unique identification of the structural shocks.

The R package **sstvars** currently accommodates three types of identification methods: recursive identification, identification by heteroskedasticity, and identification by non-Gaussianity. Identification by non-Gaussianity requires mutually independent shocks at most one of which can be Gaussian (Virolainen 2025a), and therefore, it is available only for model with independent Student's  $t$  errors distribution. In that case, the shocks are statistically identified without further assumptions, and thus we have excluded the availability of the other two identification methods to these models. Overidentifying restrictions on the impact responses of the variables to the shocks can, however, be imposed. Structural models incorporating Gaussian or conventional  $t$  shocks can be identified in **sstvars** recursively or by heteroskedasticity.

### 3.1. Recursive identification

A conventional way of identifying the shocks is to impose restrictions on the impact responses of variables. A commonly applied identification is to assume a recursive lower-triangular structure on  $B_t$ , implying that  $B_t$  is obtained as the Cholesky decomposition of the conditional covariance matrix  $\Omega_{y,t}$ . The recursive identification is straightforward to apply and it allows many of the impact responses to vary in time but constraints many of them to zero. This is particularly disadvantageous if the shock of interest is ordered last or almost last (which is typically the case in small-scale monetary policy shock applications), as then the impact responses of many of the variables to the shock of interest are zero, and therefore, time-invariant.

With our specification of the STVAR model, recursive identification does not, however, allow to impose over-identifying restrictions on the impact matrix  $B_t$ . This is because there does not exist a direct parametrization of a lower-triangular  $B_t$  such that the conditional covariance matrix  $\Omega_{y,t}$  is a weighted sum of the regime-specific covariance matrices with time-varying weights.

### 3.2. Identification via heteroskedasticity

An alternative identification method proposed by [Lütkepohl and Netšunajev \(2017\)](#) for structural VARs with smooth transitions in variances (see also the seminal paper by [Rigobon 2003](#)) identifies the shocks by simultaneously diagonalizing the covariance matrices  $\Omega_1, \dots, \Omega_M$ . This restricts the relative impact responses of the variables to be constant over time (for each shock) but does not necessarily require any zero restrictions. Since [Lütkepohl and Netšunajev \(2017\)](#) assume only two regimes and do not normalize the conditional covariance matrix of the structural error to a constant, their specification does not directly apply to our model. Therefore, we adopt the more suitable specification of [Virolainen \(2025b\)](#), and decompose the covariance matrices as

$$\Omega_m = W\Lambda_m W', \quad m = 1, \dots, M, \quad (27)$$

where the diagonal of  $\Lambda_m = \text{diag}(\lambda_{m1}, \dots, \lambda_{md})$ ,  $\lambda_{mi} > 0$  ( $i = 1, \dots, d$ ), contains the eigenvalues of the matrix  $\Omega_m \Omega_1^{-1}$  and the columns of the nonsingular  $W$  are the related eigenvectors (that are the same for all  $m$  by construction). When  $M = 2$ , the decomposition (27) always exists ([Muirhead 1982](#), Theorem A9.9), but for  $M > 2$  its existence requires that the matrices  $\Omega_m \Omega_1^{-1}$  share the common eigenvectors in  $W$ . This is, however, testable.

The impact matrix is then obtained as

$$B_t = W \left( \sum_{m=1}^M \alpha_{m,t} \Lambda_m \right)^{1/2}, \quad (28)$$

where  $\Lambda_1 = I_d$ . The shocks are identified up to ordering and sign if none of the pairs of  $\lambda_{mi}$ ,  $i = 1, \dots, d$ , is identical for all  $m = 2, \dots, M$ . Assuming that this condition is satisfied, the shocks can be labelled according to the unrestricted impact responses on  $B_t$ , and if necessary, further economically motivated restrictions can be imposed. The additional economic restrictions on the impact matrix are testable, as they are overidentifying. See [Virolainen \(2025b\)](#) for a more detailed discussion on the identification and labelling of the shocks.

Shocks identified by heteroskedasticity impose constant relative impact responses for the variables, making them unsuitable for some applications concerned with time-varying impulse response functions. Similarly to the recursive identification, this method is straightforward to apply, but unlike the recursive identification, it does not necessarily require any zero constraints on the impact responses. Compared to the recursive identification, identification by heteroskedasticity is therefore particularly advantageous when the recursive identification would imply that the shock of interest is order last. In this case, the assumption of time-invariant relative impact responses is less restrictive than the zero restrictions of the recursive identification. In contrast,



the recursive identification is particularly appealing when the shock of interest is ordered first (or almost first), as then the impact responses to the shock of interest can vary freely in time.

### 3.3. Identification by non-Gaussianity

When  $t$  shocks are mutually independent and at most one of them is Gaussian (i.e., under independent  $t$  shocks or skewed  $t$  shocks, see Sections 2.3.3 and 2.3.4), they are generally statistically identified without imposing further restrictions on the models (Virolainen 2025a). Specifically, the required identifying assumptions are:

- Assumption 1**
1. *The structural error process  $e_t = (e_{1t}, \dots, e_{dt})$  is a sequence of independent and identically distributed random vectors such that  $e_t$  is independent of  $\mathcal{F}_{t-1}$  and with each component  $e_{it}$ ,  $i = 1, \dots, d$ , having zero mean and unit variance.*
  2. *The components of  $e_t = (e_{1t}, \dots, e_{dt})$  are (mutually) independent and at most one them has a Gaussian marginal distribution.*

The first two assumptions state the requirements of the structural shocks. The latter two assumptions state that either the transition weights of the regimes take the value one or they exhibit certain small degree of variation. These are not restrictive however, since generally when the regime-switches are not discrete, there is sufficient amount of variation in the transition weights.

Under Assumption 1 the impact matrix 12 is uniquely identified at each  $t$  up to ordering and signs of its columns (Virolainen 2025a, Lemma 2). If the impact matrix is time-invariant as in Lanne, Meitz, and Saikkonen (2017), i.e.,  $B_{y,t} = B$  for some constant matrix  $B$ , it follows that the structural shocks are identified up to ordering and sign. Such statistically identified shocks can then be labelled by economic shocks based on external information. However, when the impact matrix varies in time, two complications arise. First, the impact matrix  $B_{y,t}$  is not a matrix of constant parameters, but a function of parameters, and it needs to be shown that the parameters in the specific functional form of  $B_{y,t}$  are identified. Second, since the identification of  $B_{y,t}$  at each  $t$  is only up to ordering and signs of its columns, its unique identification requires that the parameters in the functional form of  $B_{y,t}$  are restricted so that it fixes the ordering and signs of  $B_{y,t}$ .

Considering the functional form of  $B_{y,t}$  in Equation (12), in order to fix the ordering and signs of the columns of  $B_{y,t}$  it then suffices to fix the ordering and signs of the columns of  $B_1, \dots, B_M$ . This requires such constraints to be imposed on  $B_1, \dots, B_M$  that under these constraints, re-ordering or changing the signs of the columns of any of  $B_m$  would lead to an impact matrix  $B_{y,t}$  that is not observationally equivalent to the original one at some  $t$ . In other words, under the constraints, changing the ordering or signs of the columns of any of  $B_m$  should lead an impact matrix  $B_{y,t}$  that, at some  $t$ , cannot be obtained by reordering or changing the signs of the columns of the original one.

A suitable strategy for fixing the ordering and signs of the columns of  $B_1, \dots, B_M$  depends on the transition weights and distributions of the shocks, as for some specifications constraints that are not restrictive in some specifications can be overidentifying in others. Conversely, constraints that are identifying in some specifications may not be enough for identification in others. Therefore, it is useful to divide the discussion on the identification in two separate cases: when the transition weights are binary (i.e.,  $\alpha_{m,t} \in \{0, 1\}$ ) and when the transition weights exhibit more variation.

#### *Identification of the shocks in TVAR models*

Suppose the transition weights are binary,  $\alpha_{m,t} \in \{0, 1\}$ , for all  $t$  and  $m = 1, \dots, M$ . Then, at every  $t$ , the process is completely in one of the regimes and the impact matrix is  $B_{y,t} = B_m$  for

the regime  $m$  with  $\alpha_{m,t} = 1$ . In particular, following Virolainen (2025a), we assume that the binary weights are either of the threshold form (see Section 2.4.5) or exogenous (nonrandom). By Lemma 2 of Virolainen (2025a), the impact matrix  $B_{y,t}$  is thereby identified up to ordering and signs of its columns in each of the regimes, implying that identification is obtained by fixing the ordering and signs of the columns of  $B_1, \dots, B_M$ . However, whether fixing the ordering or signs of the columns of  $B_m$ ,  $m = 1, \dots, M$ , separately in each regime is overidentifying depends on the distributions of the shocks  $e_t = (e_{1t}, \dots, e_{dt})$  (Virolainen 2025a, see the discussion in)

The required identification conditions depend on the distributions of the shocks, so it is useful to make some assumptions on them. Virolainen (2025a) assumes the shocks follow independent skewed  $t$ -distributions, but the identification result extends to other suitable distributions as well, such the independent Student's  $t$ -distribution (independent  $t$ -distribution is obtained from the skewed  $t$ -distribution as the special case  $\lambda_i = 0$ ). Assuming the shocks follow independent skewed  $t$ -distributions, the identification result of Virolainen (2025a, Proposition 1) states:

**Proposition 1** *Consider the STVAR model defined in Equations (1), (2), and (12) with  $B_1, \dots, B_M$  invertible; Assumption 1 and the conditions of Lemma 2 of Virolainen (2025a) satisfied;  $\alpha_{m,t}$  either of the threshold form (26) or exogenous (nonrandom) with  $\alpha_{m,t} \in \{0, 1\}$  for all  $t$  and  $m = 1, \dots, M$ ; and the shocks  $e_{it}$ ,  $i = 1, \dots, d$ , following the skewed  $t$ -distribution with the parameters  $\nu_i > 2$  and  $\lambda_i \in (-1, 1)$  controlling kurtosis and skewness, respectively. Suppose the observed time series is  $y_{-p+1}, \dots, y_0, y_1, \dots, y_T$ , and assume the following conditions hold:*

1. *the ordering and signs of the columns of  $B_1$  are fixed,*
2.  *$(\nu_i, \lambda_i) \neq (\nu_j, \lambda_j)$  for all  $i \neq j \in \{1, \dots, d\}$ ,*
3.  *$\lambda_i \neq 0$  for all  $i = 1, \dots, d$ .*

*Then, the matrices  $B_1, \dots, B_M$  are uniquely identified almost everywhere in  $(y_{-p+1}, \dots, y_0, y_1, \dots, y_T) \in \mathbb{R}^{d(T+p)}$ .*

Besides Assumption 1 and restricting the result to certain weights functions, the remaining conditions of Lemma 2 of Virolainen (2025a) are very mild, merely stating that the AR parameters of different regimes are not all equal to each other, and there at least small degree of variation in the transition weights (for binary weights, the assumption is that there is at least one observation from each regime).

Condition 1 of Proposition 1 fixes the ordering and signs of  $B_1$ , which effectively fixes the ordering and signs of the columns of  $B_2, \dots, B_M$  under Conditions 2 and 3, respectively. Any fixed ordering and signs of the columns of  $B_1$  can be chosen without loss of generality (e.g., by making the first nonzero entry in each column is positive and ordering these entries in decreasing magnitude, given that none of them are equal). Similarly, if Condition 3 fails for some  $i \in \{1, \dots, d\}$  (meaning zero skewness), the signs of the corresponding columns of  $B_2, \dots, B_M$  can be arbitrarily fixed without loss of generality. If independent  $t$ -distribution is considered, i.e.,  $\lambda_i = 0$  for all  $i = 1, \dots, d$ , just replace Condition 3 with the condition that the signs of the columns of also  $B_2, \dots, B_M$  are fixed (any signs can be assumed without loss of generality). See Virolainen (2025a) for further details of Proposition 1.

If Condition 2 or 3 is close to break, identification is weak with respect to the ordering or signs of the corresponding columns of  $B_2, \dots, B_M$ , respectively. Somewhat surprisingly, our experience shows that even when Conditions 2 and 3 hold clearly, identification appears to be often weak with respect to the ordering and signs of the columns of  $B_2, \dots, B_M$ , at least if the parameters are estimated by the method of maximum likelihood or penalized maximum likelihood (which are the implemented methods in this package and those considered in Virolainen 2025a). Specifically, there are frequently a number of local maxima of the (penalized) log-likelihood function



with (penalized) log-likelihoods close to each other, corresponding to otherwise largely similar parameter estimates but with different the ordering and signs of the columns of  $B_2, \dots, B_M$ . As discussed in Section 3.3.3 (see also the related discussion in Virolainen 2025a), this weak identification should be appropriately dealt with when labeling the shocks, for instance, by combining Proposition 1 with additional identifying information.

### *Identification of the shocks in STVAR models*

Having established the identification result for TVAR models, or more generally to models incorporating discrete regime switches, we now consider "smoothly varying" transition weights that may take other values as well. In particular, following Virolainen (2025a), we assume that the weights are logistic or exogenous (nonrandom). The identification result can be extended to other suitable weight functions as well, but the identification of the AR and weight function parameters, and thus the identification of  $B_1, \dots, B_M$ , is shown only for these weight functions in Virolainen (2025a). The following proposition, Proposition 2 of Virolainen (2025a), states the identification result:

**Proposition 2** *Consider the STVAR model defined in Equations (1), (2), and (12); with Assumption 1 and the conditions of Lemma 2 of Virolainen (2025a) satisfied; and let the transition weights be either exogenous (nonrandom) or follow the logistic process defined in (19) (with  $M = 2$  assumed). Suppose the observed time series is  $y_{-p+1}, \dots, y_0, y_1, \dots, y_T$ , and assume the following conditions hold:*

1. *the ordering and signs of the columns of  $B_1$  are fixed, and*
2. *the vector  $(\alpha_{1,t}, \dots, \alpha_{M,t})$  takes a value for some  $t$  such that none of its entries is zero.*

*Then, for models with logistic transition weights, the matrices  $B_1, \dots, B_M$  are uniquely identified almost everywhere in  $[B_1 : \dots : B_M] \in \mathbb{R}^{d \times dM}$ ,  $(y_{-p+1}, \dots, y_0, y_1, \dots, y_{T-1}) \in \mathbb{R}^{d(T+p-1)}$  and, if the switching variable  $z_t$  is exogenous,  $(z_1, \dots, z_T) \in \mathbb{R}^T$ . For models with nonrandom exogenous transition weights, this result holds almost everywhere in  $[B_1 : \dots : B_M] \in \mathbb{R}^{d \times dM}$ .*

Condition 1 of Proposition 2 states that the ordering and signs of the columns of  $B_1$  should be fixed, which fixes the ordering and signs of the columns of  $B_{y,t}$ . If the shocks follow the skewed  $t$ -distributions (or independent  $t$ -distributions), any fixed ordering and signs of the columns of  $B_1$  can be assumed without loss of generality, as reordering the columns would just reorder the shocks and swapping a sign of a column corresponds to swapping the sign of the related shock and skewness parameter. Condition 2 states that the transition weights should, for some  $t$ , be strictly positive for all of the regimes. Similarly to the structural TVAR model discussed in Section 3.3.1, our experience shows that identification appears to be often weak with respect to the ordering and signs of the columns of  $B_2, \dots, B_M$ , which should be appropriately dealt with when labelling the shocks, as discussed in the next section.

### *Labelling the shocks and blended identification*

Similarly to the identification by heteroskedasticity, the shocks are statistically identified without further restrictions or assumptions. But labelling the shocks by the economic shocks of interest requires external information. Also, labelling the shocks based on the estimates of the impact matrices  $B_1, \dots, B_M$  might not always be straightforward, because the same shock needs to be associated with the same column of the impact matrix  $B_m$  in all regimes. For example, if a positive supply shock increases output and decreases prices on impact, and it should do so in all regimes, labelling the  $i$ th shock as the supply shock requires the  $i$ th column of all  $B_1, \dots, B_M$  to satisfy such signs. However, as discussed in Sections 3.3.1 and 3.3.2, identification

can be weak with respect to the ordering and signs of the columns of  $B_2, \dots, B_M$ , which should be appropriately accounted for when labelling the shocks.

Assuming that the parameters are estimated by the method of maximum likelihood (ML) or penalized maximum likelihood (PML) described in Section 4, weak identification often results in multiple local maxima of the (penalized) log-likelihood function with the related (penalized) log-likelihoods close to each other. If identification is weak with respect to the ordering and signs of the columns of  $B_2, \dots, B_M$ , these local solutions often correspond to otherwise largely similar parameter estimates but with different ordering and signs of the columns of  $B_2, \dots, B_M$ . In **sstvars**, different local solutions can be conveniently examined with the functions **alt\_stvar** and **filter\_estimates** by adjust their arguments **which\_largest**.

To formally obtain the local solution that facilitates labelling the shock of interest to the same column of the impact matrix in all regimes, we recommend employing a "blended identification" strategy that combines identification by non-Gaussianity with additional identifying information (cf. [Carriero, Marcellino, and Tornese 2024](#)). As a concrete example, if a supply shock should increase output and decrease prices at impact in all regimes, selecting the best local solution where the same column in all  $B_1, \dots, B_M$  satisfies such signs amounts to combining identification by non-Gaussianity with the appropriate short-run sign restrictions. In addition to short-run sign restrictions, also other types of information can be exploited in blended identification, including narrative restrictions and testable zero impact effect restrictions, for instance.

The ML and PML estimation methods implemented to **sstvars** produce a set of local solutions to the estimation problem, so the researcher can check whether there are multiple local solutions that yield log-likelihoods close to each other. Given that such local solutions are otherwise largely similar but incorporating different ordering or signs of some of the columns of  $B_m$  in some regime, the weak identification is related to the ordering and signs of the columns. In practice, one can thereby simply choose among the almost equally well fitting local solutions the solution that satisfies the additional constraints imposed in the "blended identification method", i.e., the local solution that facilitates labelling the shock(s) of interest to the same column of the impact matrix in all of the regimes. In **sstvars** different local solutions are most conveniently examined with the functions **alt\_stvar** and **filter\_estimates**. Structural models that impose overidentifying short-run sign or zero restrictions can, in turn, be estimated with the function **fitSSTVAR**.

## 4. Estimation

We consider maximum likelihood estimation (ML) as well as penalized maximum likelihood (PML) estimation of the STVAR models. The former maximizes the log-likelihood function, and the latter maximizes the penalized log-likelihood function that penalizes from estimates that are close to the boundary or outside the stability region (i.e., a region where the usual stability condition is satisfied by each of the regimes). In either case, the numerical estimation will be conducted in two or three phases, which are discussed after introducing the log-likelihood function and the penalized log-likelihood function.

The parameters of the reduced form STVAR model are collected to the vector

$$\boldsymbol{\theta} = (\phi_1, \dots, \phi_M, \varphi_1, \dots, \varphi_M, \sigma, \alpha, \nu), \quad (29)$$

where  $\varphi_m = (\text{vec}(A_{m,1}), \dots, \text{vec}(A_{m,p}))$ ,  $m = 1, \dots, M$ ,  $\sigma$  contains the covariance matrix parameters ( $(\text{vech}(\Omega_1), \dots, \text{vech}(\Omega_M))$  for Gaussian and Student  $t$  models, and  $(\text{vec}(B_1), \dots, \text{vec}(B_M))$  for independent Student's  $t$  models),  $\alpha$  contains the transition weight parameters,  $\nu$  contains the distribution degrees of freedom and skewness parameter(s) (if any, completely omitted for Gaussian models),  $\text{vec}$  is a vectorization operator that stacks the columns of a matrix on

top of each other, and `vech` stacks the columns of a matrix from the main diagonal downwards (including the main diagonal). Structural models identified by heteroskedasticity assume  $\sigma = (\text{vec}(W), \lambda_m, \dots, \lambda_M)$ ,  $\lambda_m = (\lambda_{m1}, \dots, \lambda_{md})$ ,  $m = 2, \dots, M$ , whereas structural models identified recursively or by non-Gaussianity use the same parameter vector as reduced form models.

If relative stationary densities are used as transition weights,  $\alpha = (\alpha_1, \dots, \alpha_{M-1})$  ( $\alpha_M$  is not included because it is obtained from the constraint  $\sum_{m=1}^M \alpha_m = 1$ ), where we assume, for identification, that  $\alpha_1, \dots, \alpha_{M-1}$  are in a decreasing order. For exponential and logistic transition weights,  $\alpha = (c, \gamma)$ , for multinomial logit transition weights  $\alpha = (\gamma_1, \dots, \gamma_{M-1})$  ( $\gamma_M$  is not included because  $\gamma_M = 0$  is assumed for identification), and with threshold transition weights,  $\alpha = (r_1, \dots, r_{M-1})$ . With exogenous transition weights, the parameter  $\alpha$  is dropped.

#### 4.1. The log-likelihood function

`sstvars` employs the method of maximum likelihood (ML) for estimating the parameters of the STVAR models. Indexing the observed data as  $y_{-p+1}, \dots, y_0, y_1, \dots, y_T$ , the conditional log-likelihood function conditional on the initial values  $\mathbf{y}_0 = (y_{-p+1}, \dots, y_0)$  is given as

$$L_t(\boldsymbol{\theta}) = \sum_{t=1}^T l_t(\boldsymbol{\theta}) = \sum_{t=1}^T \log d_d(y_t; \mu_{y,t}, \Omega_{y,t}, \nu). \quad (30)$$

where  $d_d(y_t; \mu_{y,t}, \Omega_{y,t}, \nu)$  is the  $d$ -dimensional conditional density of the process, conditional on  $\mathcal{F}_{t-1}$ , at time  $t$ , given in Section 2.3. When the conditional distribution is Gaussian, the degrees of freedom parameter(s)  $\nu$  is/are dropped from the right side of (30). The ML estimator of  $\boldsymbol{\theta}$  maximizes the log-likelihood function  $L_t(\boldsymbol{\theta})$  over the parameter space specified in the below assumption.

We summarize the constraints (often) imposed on the parameter space in the following assumption.

**Assumption 2** *The true parameter value  $\boldsymbol{\theta}_0$  is an interior point of  $\boldsymbol{\Theta}$ , which is a compact subset of  $\{\boldsymbol{\theta} = (\phi_1, \dots, \phi_M, \varphi_1, \dots, \varphi_M, \sigma, \alpha, \nu) \in \mathbb{R}^{M(d+d^2p+d(d+1)/2)} \times S \times V: \Omega_m \text{ is positive definite (or } B_m \text{ is invertible) for all } m = 1, \dots, M, \text{ and Condition 1 holds.}\}$ .*

Above,  $S = (0, 1)^{M-1}$  for relative densities transition weights,  $S = \mathbb{R} \times (0, \infty)$  for logistic and exponential weights,  $S = \mathbb{R}^{(M-1)k}$  for multinomial logistic weights,  $S = \mathbb{R}^{M-1}$  for threshold weights, and  $S$  is an empty set for exogenous weights. Also,  $V$  is omitted for Gaussian models,  $V = (2, \infty)$  for Student's  $t$  models,  $V = (0, \infty)^d$  for independent Student's  $t$  models, and  $V = (0, \infty)^d \times (0, 1)^d$  for skewed  $t$  models.

As noted before, Condition 1 is not necessary, but it ensures stationarity and ergodicity of the process, unless the transition weights are exogenous. In estimation, we impose the more easily verified Condition 2, however, and the sufficient condition can be check after the estimation with the function `bound_JSR`. As noted in Section 2.2, the sufficient condition is in practice usually satisfied if the necessary condition is satisfied and not very close to being violated. Given that under Condition 1 the process is ergodic stationary, there is no particular reason to believe that the standard asymptotic results of consistency and limiting Gaussian distribution would not apply to the ML estimator.

As also noted before, we also allow estimation does not impose Condition 2, in particular, with the penalized ML estimation. This is particularly beneficial when the data is persistent or the estimated model is large, as allowing the estimation algorithm to explore the parameter space more freely generally seems to improve the performance of the estimation algorithms (in our experience). Moreover, in the three-step estimation the initial (nonlinear) LS estimates do not often satisfy the usual stability condition, and allowing for unstable estimates thereby allows to

use them more effectively. When penalized ML estimation is used, instability of the estimates is penalized in the objective function, so that the optimization algorithm often eventually converge back to the stability region.

## 4.2. The penalized log-likelihood function

The penalized likelihood function can be written as

$$PL(\boldsymbol{\theta}) = L(\boldsymbol{\theta}) - P(\boldsymbol{\theta}), \quad (31)$$

where  $L(\boldsymbol{\theta})$  is defined in (30) and  $p(\boldsymbol{\theta}) \geq 0$  is a penalization term that penalizes the log-likelihood function from parameter values that are close to entering or are in an uninteresting region of the parameter space. We focus on avoiding estimates that do not satisfy the usual stability condition for all the regimes. Hence, we define the penalization term as

$$P(\boldsymbol{\theta}) = \kappa T d \sum_{m=1}^M \sum_{i=1}^{dp} \max\{0, |\rho(\mathbf{A}_m(\boldsymbol{\theta}))_i| - (1 - \eta)\}^2, \quad (32)$$

where  $|\rho(\mathbf{A}_m(\boldsymbol{\theta}))_i|$  is the modulus of the  $i$ th eigenvalue of the companion form AR matrix of Regime  $m$ , the tuning parameter  $\eta \in (0, 1)$  determines how close to the boundary of the stability region the penalization starts, and  $\kappa > 0$  determines the strength of the penalization. The PML estimate of  $\boldsymbol{\theta}$  maximizes the log-likelihood function (31) with respect to  $\boldsymbol{\theta}$ .

Whenever the companion form AR matrix of a regime has eigenvalues greater than  $(1 - \eta)$  in modulus, the penalization term (32) is greater than zero, and it increases in the modulus of these eigenvalues. Finally, our penalization term has the multiplicative coefficient  $Td$ , as the value of the log-likelihood function (30) tends to scale both with number of observations  $T$  and with the number of variables  $d$ . The default values of tuning parameter values are  $\eta = 0.05$  and  $\kappa = 0.2$ , thus, penalizing significantly estimates that are outside the stability region, while maintaining some flexibility near the boundary. In `sstvars`, the tuning parameter values can be adjusted with the argument `penalty_params`.

Penalized log-likelihood estimation can be used both with the two-phase and three-step estimation procedures. As already noted, the penalized ML estimation is particularly useful with persistent data and more complex models, because it allows the estimation algorithm to explore the parameter space more freely, assuming that `allow_unstab=TRUE` is used. The penalized ML estimation is particularly useful in the three-step estimation when the initial (nonlinear) LS estimates are unstable, as the penalization term in the objective function often eventually drives the optimization algorithm back to the stability region.

## 4.3. Two-phase estimation procedure

Finding the ML estimate amounts maximizing the log-likelihood function (30) over a high dimensional parameter space satisfying the constraints summarized in Assumption 2. Due to the complexity of the log-likelihood function, numerical optimization methods are required. The maximization problem can be challenging in practice due to the dependence of the transition weights on the preceding observations, which induces a large number of modes to the surface of the log-likelihood function, and large areas to the parameter space, where it is flat in multiple directions.

Therefore, we follow Meitz *et al.* (2023) and Virolainen (2022b) and employ a two-phase estimation procedure that is run for a large number of times. In the first phase, a genetic algorithm is used to find parameter values (hopefully) near local maximums. Since genetic algorithms tend to converge slowly near local solutions, a gradient based variable algorithm (Nash 1990, algorithm 21, implemented by R Core Team 2022) is ran for each of the starting values, resulting

in a number of alternative local solutions. Some of the estimation rounds may end up in saddle points or near-the-boundary points that are not local solutions, and some of the local solutions may be inappropriate for econometric inference (for instance, there might be only a few observations from some of the regimes). After the estimation rounds have been ran, the researcher can choose the local solution that maximizes the log-likelihood among the appropriate local solutions. Inappropriate solutions are automatically filtered by **sstvars**, but this functionality can also be turned off and the researchers can use estimates based any estimation round. The R package **sstvars** employs a modified genetic algorithm that works similarly to the one described in the R packages **uGMAR** (Virolainen 2018b) and **gmvarKit** (Virolainen 2018a) (the genetic algorithm and implemented in former is briefly described in Virolainen 2022b). See Virolainen (2022a, Chapter 3) for a related discussion on complex numerical estimation problems using the two-phase procedure.

#### 4.4. Three-step estimation procedure

The two-phase procedure facilitates finding the ML estimate, but often a very large number of estimation rounds is required to obtain reliable results, particularly when the model is large. Running a very large number of estimation rounds, however, often requires very substantial computation power, making the estimation tedious in practice. Consequently, estimation of large models often infeasible with computer available for the practitioners.

To address the issue of the extreme computation power required for estimating large models, Virolainen (2025a) propose a more computationally efficient three-step estimation procedure by adapting the three-step procedure of Lanne *et al.* (2017) to nonlinear SVAR models. The three-step estimation procedure operates with the following steps:

1. Estimate the autoregressive and the threshold parameters  $\theta_1 \equiv (\phi_1, \dots, \phi_M, \varphi_1, \dots, \varphi_M, \alpha)$  by the method of (penalized) nonlinear least squares NLS. Denote this estimate of  $\theta_1$  as  $\hat{\theta}_1^{NLS}$ .
2. Estimate the error distribution parameters  $\theta_2 \equiv (\sigma, \nu)$  by (P)ML with a genetic algorithm conditionally on the PLS estimate  $\hat{\theta}_1^{NLS}$ . Denote the obtained estimate of  $\theta_2$  as  $\hat{\theta}_2^{GA}$ .
3. Estimate the full parameter vector  $\theta = (\theta_1, \theta_2)$  by (P)ML by initializing a standard gradient based optimization algorithm from  $(\hat{\theta}_1^{NLS}, \hat{\theta}_2^{GA})$ .

Step 1 obtains initial estimates for the AR and weight function parameters, and it is comparable to Step 1 of Lanne *et al.* (2017) but with the nonlinearity of the model explicitly accounted for. Notably, the NLS estimation often produces estimates that do not satisfy the usual stability condition for each of the regimes, and thus allowing for instability (but penalizing it) facilitates utilization of the NLS estimates in our three-step procedure. Step 2 is comparable to Step 2 of Lanne *et al.* (2017), but the multimodality of the (penalized) log-likelihood function is taken into account by making use of a robust estimation algorithm that is able to escape from local maxima. Step 3 finalizes the estimation by initializing a standard gradient based optimization algorithm from the initial estimates obtained from the previous steps, often converging to a nearby local maximum of the (penalized) log-likelihood function. **sstvars** uses the variable metric algorithm described in (Nash 1990, Algorithm 21), which is implemented to the standard R function **optim**, but other gradient based algorithms could be also used. This steps is comparable to Step 3 estimation of Lanne *et al.* (2017).

Due to the complexity and high multimodality of the log-likelihood function, Step 2 should be ran a number times to improve the reliability of the results. Since multiple estimation rounds are ran, the procedure produces a set of estimates, possibly corresponding to different local maximums of the (penalized) log-likelihood function. However, due to the initial estimates



obtained from Step 1, a smaller number of estimation rounds is required than in the two-phase procedure.

#### 4.5. Examples of unconstrained estimation

In this section, we demonstrate how to estimate STVAR models with `sstvars`. **The examples use two-phase estimation, but three-step estimation can be employed in a similar manner.** In the examples, we only consider  $p = 1$  models for simplicity and merely because then the code outputs fit in the margins better. This order may not be the best in the modeling perspective, however.

In `sstvars`, the STVAR models are defined as class `stvar` S3 objects, which can be created with given parameter values using the constructor function `STVAR` (see Section 10) or by using the estimation function `fitSTVAR`, which estimates the parameters and then builds the (reduced form) model. Structural models are estimated based on a reduced form model with the function `fitSSTVAR`. For estimation, `fitSTVAR` needs to be supplied with a multivariate time series and the arguments specifying the model. The necessary arguments for specifying the model include the autoregressive order `p`, the number of regimes `M`, the transition weight function `weight_function`, with some weight functions the switching variable(s) `weightfun_pars`, and the conditional distribution `cond_dist`.

Additional arguments may be supplied to `fitSTVAR` in order to specify, most importantly, how many estimation rounds should be performed (`nrounds`) and how many central processing unit (CPU) cores should be used in the estimation (`ncores`). Some of the estimation rounds may end up in local-only maximum points or saddle points, but reliability of the estimation results can be improved by increasing the number of estimation rounds. A large number of estimation rounds may be required particularly when the number of regimes is large or there are many variables in the data, as the surface of the log-likelihood function becomes increasingly more challenging. It is also possible to adjust the settings of the genetic algorithm that is used to find the starting values. The available options are listed in the documentation of the function `GAFit` to which the arguments adjusting the settings will be passed.

**In general, we recommend being conservative with choice of  $M$  due to the identification problems induced if the number of regimes is chosen too large. Also, estimation of models that contain more than two regimes can be extremely challenging. Another important thing to know about estimation is that the estimation algorithm performs very poorly if some of the AR coefficients are very large, substantially larger than one. This means that you need to scale each component time series so that they vary approximately in the same magnitude. For instance, typically in macroeconomic time series, log-differences should be multiplied by hundred. If the suitable scales are not obvious, you can try out different scales and estimate linear VARs with your favorite package to see whether the AR coefficients are in a reasonable range. When a suitable scale is found, proceed to the STVAR models.**

We illustrate the use of `sstvars` with a quarterly series consisting of two U.S. variables: the percentage change of real GDP and the percentage change of GDP implicit price deflator, covering the period from 1959Q1 to 2019Q4. The following code fits a  $STVAR(p = 1, M = 2)$  model with Student's  $t$  conditional distribution and logistic transition weights by performing 24 estimation rounds with 8 CPU cores. **In practice, hundreds or even thousands of estimation rounds is often required to obtain reliable results. The larger the dimension of the series is and the larger the order of the model is and the more there are regimes, the more estimation rounds is required. The model in our example is easy to estimate, as it is small in dimension and order.** We set the switching variable to be the first lag of the second variable, i.e., the GDP deflator by setting argument `weightfun_pars=c(2, 1)`.

The argument `seeds` supplies the seeds that initialize the random number generator at the beginning of each call to the genetic algorithm, thereby yielding reproducible results.

```
R> library(sstvars)
R> data("gdpdef", package="sstvars")
R> fit12 <- fitSTVAR(gdpdef, p=1, M=2, weight_function="logistic",
+   weightfun_pars=c(2, 1), cond_dist="Student", estim_method="two-phase",
+   nrounds=24, ncores=8, seeds=1:24)
```

Using 8 cores for 24 estimations rounds...

PHASE 1: Estimating all the parameters with a genetic algorithm...

```
|+++++| 100% elapsed=16s
```

Results from the genetic algorithm:

The lowest loglik: -278.505

The largest loglik: -251.32

PHASE 2: Estimating all the parameters with a variable metric algorithm...

```
|+++++| 100% elapsed=02s
```

Results from the variable metric algorithm:

The lowest loglik: -273.766

The largest loglik: -250.236

Filtering inappropriate estimates...

Filtered through 0 inappropriate estimates with a larger log-likelihood

Calculating approximate standard errors...

Finished!

The progression of the estimation process is reported with a progress bar giving an estimate of the remaining estimation time. Also statistics on the spread of the log-likelihoods are printed after each estimation phase. The progress bars are generated during parallel computing with the package **pbapply** (Solymos and Zawadzki 2020).

Because the log-likelihood function is highly multimodal, and the estimation algorithm is ran a large number of times, it produces a set of local solutions, possibly representing various modes in the log-likelihood function. Some of the local solutions may be inappropriate for econometric inference, for instance, because they contain a near-singular error term covariance matrix or regimes that have only a very small number of observations generated (even partially) from them. Such inappropriate solutions, i.e., estimates that are not solutions of interest, are filtered automatically by **sstvars**. Specifically, solutions that incorporate a near-singular error term covariance matrix (any eigenvalue less than 0.002), any modulus "bold A" eigenvalues larger than 0.9985 (indicating the necessary condition for stationarity is close to break), or transition weights such that they are close to zero for almost all  $t$  for at least one regime. With relative densities transition weights, also solutions in which a weight parameter estimate is close to zero are filtered out.

The various local solutions (from any estimation round) can also be easily browsed and examined by using the function `alt_stvar` and adjusting its argument `which_largest` or `which_round`. Also, the function `filter_estimates` can be used to conveniently browse various estimates that are not deemed inappropriate.

The estimates can be examined with the `print` method.

```
R> print(fit12)
```

```
logistic Student STVAR model, reduced form model no AR_constraints,
no mean_constraints,
```

```
p = 1, M = 2, d = 2, #parameters = 21, #observations = 243 x 2
Switching variable: GDPDEF with lag 1.
```

```
Regime 1
Degrees of freedom: 7.70 (for all regimes)
Regime means: 0.71, 0.49
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 0.63 ] + [ 0.35 -0.35 ] y1.1 + [ 0.37 0.00 ]      eps1
2 y2   [ 0.14 ]   [ 0.06 0.62 ] y2.1   [ 0.00 0.03 ]      eps2
```

```
Regime 2
Weight params: 1.22 (location), 5.01 (scale)
Regime means: 0.77, 1.76
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 2.41 ] + [ 0.13 -0.99 ] y1.1 + [ 1.29 -0.06 ]      eps1
2 y2   [ 0.67 ]   [-0.04 0.64 ] y2.1   [-0.06 0.19 ]      eps2
```

The parameter estimates are reported for each mixture component separately so that the estimates can be easily interpreted. Each regime's autoregressive formula is presented in the form

$$y_t = \varphi_{m,0} + A_{m,1}y_{t-1} + \dots + A_{m,p}y_{t-p} + \Omega_m^{1/2}e_t. \quad (33)$$

The other statistics are listed above the formula: the degrees of freedom parameter estimate, the unconditional means of the regimes, and estimates of the transition weight parameters. For models incorporating mutually independent Student's  $t$  shocks (`cond_dist="ind_Student"`), estimates of the impact matrices  $B_m$ ,  $m = 1, \dots, M$ , are presented in the place of the covariance matrices  $\Omega_m$ . Based on the estimate of the location parameter and the conditional means in the above printout, the second regime accommodates periods of high inflation and the first regime thereby periods of lower inflation.

A more detailed printout is obtained with the `summary` method as follows:

```
R> summary(fit12)
```

```
logistic Student STVAR model, reduced form model, no AR_constraints,
no mean_constraints,
```

```
p = 1, M = 2, d = 2, #parameters = 21, #observations = 243 x 2
Switching variable: GDPDEF with lag 1.
```

```
loglik/T: -1.03, AIC: 2.23, HQIC: 2.35, BIC: 2.53
```

```
Regime 1
Degrees of freedom: 7.70 (for all regimes)
Moduli of 'bold A' eigenvalues: 0.49, 0.49
Cov. matrix 'Omega' eigenvalues: 0.37, 0.03
Regime means: 0.71, 0.49
Regime sdevs: 0.66, 0.24
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 0.63 ] + [ 0.35 -0.35 ] y1.1 + [ 0.37 0.00 ]      eps1
2 y2   [ 0.14 ]   [ 0.06 0.62 ] y2.1   [ 0.00 0.03 ]      eps2
```



Error term correlation matrix:

```
      [,1] [,2]
[1,] 1.000 0.028
[2,] 0.028 1.000
```

Regime 2

```
Moduli of 'bold A' eigenvalues: 0.71, 0.06
Cov. matrix 'Omega' eigenvalues: 1.29, 0.18
Weight params: 1.22 (location), 5.01 (scale)
Regime means: 0.77, 1.76
Regime sdevs: 1.32, 0.59
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 2.41 ] + [ 0.13 -0.99 ] y1.1 + [ 1.29 -0.06 ] eps1
2 y2   [ 0.67 ]   [ -0.04  0.64 ] y2.1   [ -0.06  0.19 ] eps2
```

Error term correlation matrix:

```
      [,1] [,2]
[1,] 1.00 -0.12
[2,] -0.12 1.00
```

Print approximate standard errors with the argument `'standard_error_print=TRUE'`.

The above summary printout shows additional information compared to the print method, including moduli of the eigenvalues of the companion form AR matrices (to assess how close Condition 2 is to break), eigenvalues of the error term covariance matrices (to assess how close they are to being non-positive definite), marginal standard deviations of the variables in each regime, error term correlation matrices, and the log-likelihood as well as the values of the information criteria divided by the number of observations  $T - p$ .

Approximate standard errors can be printed by specifying the argument `standard_error_print = TRUE` in the print method, which prints the standard errors in the same form as the print method prints the estimates. There is no standard error for the intercepts if mean parametrization is used (by setting `parametrization = "mean"` in `fitSTVAR`) and vice versa. In order to obtain standard errors for the regimewise unconditional means or intercepts, one can easily swap between the mean and intercept parametrizations with the function `swap_parametrization`. Note that approximate standard errors are based on the mere assumption of the standard Gaussian asymptotic distribution of the estimator. Hence, they should be interpreted with caution. Finally, we show how STVAR models incorporating exogenous transition weights can be estimated. In the below example, we assume two regimes ( $M = 2$ ), autoregressive order one ( $p = 1$ ), and mutually independent Student's  $t$  shocks. As an arbitrary example of exogenous transition weights, we draw the weights of Regime 1 by random from the uniform distribution, and construct the transition weights matrix as follows:

```
R> set.seed(1)
R> tw1 <- runif(nrow(gdpdef) - 1) # Transition weights of Regime 1
R> twmat <- cbind(tw1, 1 - tw1) # Transition weights of both regimes
```

Each column of the transition weight matrix gives the weights of the corresponding regime, and each row gives the time  $t$  weights. There should be as many rows as there are observations in the data minus the autoregressive order  $p$ , as the first  $p$  observations are the initial values. Each row should sum to one and all the elements need to be weakly larger than zero.

A two-regime  $p = 1$  STVAR model with the randomly drawn exogenous transition weights can be estimated by specifying the transition weights in the argument `weightfun_pars` as follows:

```
R> fitexo12 <- fitSTVAR(gdpdef, p=1, M=2, weight_function="exogenous",
+   weightfun_pars=twmat, cond_dist="ind_Student", estim_method="two-phase",
+   nrounds=24, ncores=8, seeds=1:24)
```

Using 8 cores for 24 estimations rounds...

PHASE 1: Estimating all the parameters with a genetic algorithm...

```
|+++++| 100% elapsed=24s
```

Results from the genetic algorithm:

The lowest loglik: -287.191

The largest loglik: -272.426

PHASE 2: Estimating all the parameters with a variable metric algorithm...

```
|+++++| 100% elapsed=01s
```

Results from the variable metric algorithm:

The lowest loglik: -271.593

The largest loglik: -271.593

Filtering inappropriate estimates...

Filtered through 0 inappropriate estimates with a larger log-likelihood

Calculating approximate standard errors...

Finished!

We examine the estimates with the summary printout:

```
R> summary(fitexo12)
```

exogenous ind\_Student STVAR model, reduced form model, no AR\_constraints,  
no mean\_constraints,

```
p = 1, M = 2, d = 2, #parameters = 22, #observations = 243 x 2
```

loglik/T: -1.12, AIC: 2.42, HQIC: 2.54, BIC: 2.73

Regime 1

Degrees of freedom: 3.73, 3.78 (for all regimes)

Moduli of 'bold A' eigenvalues: 0.90, 0.08

Cov. matrix 'Omega' eigenvalues: 0.51, 0.10

Regime means: 0.68, 0.47

Regime sdevs: 0.72, 0.70

```
      Y      phi0      A1      B
1 y1 = [ 0.63 ] + [ 0.08 -0.01 ] y1.1 + [ 0.71 0.11 ] eps1
2 y2   [ 0.03 ]   [ 0.02 0.90 ] y2.1   [ 0.03 -0.31 ] eps2
```

Error term correlation matrix:

```
      [,1] [,2]
[1,] 1.000 -0.065
[2,] -0.065 1.000
```

Regime 2

Moduli of 'bold A' eigenvalues: 0.87, 0.48

Cov. matrix 'Omega' eigenvalues: 0.82, 0.06

Regime means: 0.90, 0.91

Regime sdevs: 1.03, 0.54

```

      Y      phi0      A1      B
1 y1 = [ 0.65 ] + [ 0.47 -0.19 ] y1.1 + [ 0.78 0.47 ] eps1
2 y2   [ 0.09 ]   [ 0.02 0.88 ] y2.1   [ 0.17 -0.20 ] eps2

```

Error term correlation matrix:

```

      [,1] [,2]
[1,] 1.00 0.16
[2,] 0.16 1.00

```

Print approximate standard errors with the argument 'standard\_error\_print=TRUE'.

As the above printout shows, the impact matrices of the regimes,  $B_1, \dots, B_M$  are directly estimated and their estimates presented in the place where the estimates of  $\Omega_m$  are for models with `cond_dist="Gaussian"` or `cond_dist="ind_Student"`.

#### 4.6. Further examination of the estimates

In addition to examining the summary printout, it is often useful to visualize the model by plotting the transition weights together with the time series. That is exactly what the `plot` method for STVAR models does. The following command creates the time series plot along with estimated transition weights:

```
R> plot(fit12, type="series")
```

The resulting plot is presented in Figure 1. The figure confirms our earlier observation: the second regime is dominant during the periods of high inflation, particularly in the volatile periods of 1970's and 1980's.

It is also sometimes interesting to examine the time series of (one-step) conditional means of the process along with the time series the model was fitted to. This can be done conveniently with the function by setting the argument `plot_type="cond_mean"` in the `plot` method. This plot depicts the contribution of each regime to the conditional mean of the process and how close the conditional mean is to the observed series in each point of time.

The variable metric algorithm employed in the final estimation does not necessarily stop at a local maximum point. The algorithm might also stop at a saddle point or near a local maximum, when the algorithm is not able to increase the log-likelihood, or at any point, when the maximum number of iterations has been reached. In the latter case, the estimation function throws a warning, but saddle points and inaccurate estimates need to be detected by the researcher.

It is well known that in a local maximum point, the gradient of the log-likelihood function is zero, and the eigenvalues of the Hessian matrix are all negative. In a local minimum, the eigenvalues of the Hessian matrix are all positive, whereas in a saddle point, some of them are positive and some negative. Nearly numerically singular Hessian matrices occur when the surface of the log-likelihood function is very flat about the estimate in some directions. This particularly happens when the transition weights  $\alpha_{m,t}$  are estimated close to zero for all  $t = 1, \dots, T$  for some regime  $m$ .

**sstvars** provides several functions for evaluating whether the estimate is a local maximum point. The function `get_foc` returns the (numerically approximated) gradient of the log-likelihood function evaluated at the estimate, and the function `get_soc` returns eigenvalues of the (numerically approximated) Hessian matrix of the log-likelihood function evaluated at the estimate.




Figure 1: The figure produced by the command `plot(fit12)`. On the top, a quarterly series consisting of two U.S. variables: the percentage change of real GDP and the percentage change of GDP implicit price deflator, covering the period from 1959Q1 to 2019Q4. On the bottom, the estimated transition weights of the STVAR model `fit12gs` fitted the series.

The numerical derivatives are calculated using a central difference approximation

$$\frac{\partial L(\boldsymbol{\theta})}{\partial \theta_i} \approx \frac{f(\boldsymbol{\theta} + \mathbf{h}^{(i)}) - f(\boldsymbol{\theta} - \mathbf{h}^{(i)})}{2h}, \quad h > 0, \quad (34)$$

where  $\theta_i$  is the  $i$ th element of  $\boldsymbol{\theta}$  and  $\mathbf{h}^{(i)} = (0, \dots, 0, h, 0, \dots, 0)$  contains  $h$  as its  $i$ th element. By default, the difference  $h = 6 \cdot 10^{-6}$  is used.

For example, the following code calculates the first order condition for the G-StMVAR model `fit12`:

```
R> get_foc(fit12)

[1] 0.0013350596 0.0338560277 0.0001781006 0.0027846292 0.0011998654
[6] 0.0313456858 0.0003378474 0.0179776013 0.0003544770 0.0051172636
[11] 0.0002186068 0.0022140706 -0.0055795827 0.0071732765 0.5804849358
[16] -0.0002884765 0.0008926041 0.0258026645 -0.0204821154 -0.0018359291
[21] 0.0004548983
```

and the following code calculates the second order condition:

```
R> get_soc(fit12)

[1] 0.0013350596 0.0338560277 0.0001781006 0.0027846292 0.0011998654
[6] 0.0313456858 0.0003378474 0.0179776013 0.0003544770 0.0051172636
[11] 0.0002186068 0.0022140706 -0.0055795827 0.0071732765 0.5804849358
[16] -0.0002884765 0.0008926041 0.0258026645 -0.0204821154 -0.0018359291
[21] 0.0004548983
```




Figure 2: The figure produced by the command `profile_logliks(fit12)`. The graphs of the profile log-likelihood functions of the logistic Student's  $t$  STVAR model drawn about the estimate. The red vertical lines denote the estimate.

All eigenvalues of the Hessian matrix are negative, which points to a local maximum, and the gradient of the log-likelihood function is close to zero. The gradient is not exactly zero, because it is based on a numerical approximation. It is also possible that the estimate is inaccurate, because it is based on approximative numerical estimation, and the estimates are therefore not expected to be exactly accurate. Whether the estimate is a local maximum point with accuracy that is reasonable enough, can be evaluated by plotting the graphs of the profile log-likelihood functions about the estimate. In `sstvars`, this can be done conveniently with the function `profile_logliks`.

The exemplify, the following command plots the graphs of profile log-likelihood functions of the estimated G-StMVAR model `fit12`:

```
R> profile_logliks(fit12)
```

The resulting plot is presented in Figure 2.

The output shows that the estimate's accuracy is reasonable, as changing any individual parameter value marginally would not increase the log-likelihood much. The argument `scale` can be adjusted to shorten or lengthen the interval shown in the horizontal axis. If one zooms in enough by setting `scale` to a very small number, it can be seen that the estimate is not exactly at the local maximum, but it is so close that moving there would not increase the log-likelihood notably. The argument `precision` can be adjusted to increase the number of points the graph is based on. For faster plotting, it can be decreased, and for more precision, it can be increased. The argument `which_pars` is used to specify the parameters whose profile log-likelihood functions should be plotted. This argument is particularly useful when creating as many plots as there are parameters in the model to a single figure would cause the individual plots to be

very small. In such a case, profile log-likelihood functions for subsets of the parameters can be plotted separately by specifying this argument accordingly.

We have discussed tools that can be utilized to evaluate whether the found estimate is a local maximum with a reasonable accuracy. It is, however, more difficult to establish that the estimate is the global maximum. With **sstvars**, the best way to increase the reliability that the found estimate is the global maximum (among the appropriate solutions), is to run more estimation rounds by adjusting the argument **nrounds** of the estimation function **fitSTVAR**.

If the model is very large, a very large number of estimation rounds may be required to find the global maximum. If there are two regimes in the model,  $p$  is reasonable, and the dimension of the time series at most four, the required number of estimation rounds typically varies from several hundred to several thousand depending on the model and the data. In the simpler models, less estimation rounds are required. In the larger models, and in particular if  $M > 2$  or  $d > 4$ , a significantly large number of estimation rounds may be required obtain the MLE. Another thing that makes the estimation more challenging, are exotic parameter constraints that do not reduce the dimension of the parameter much. Constraints that greatly reduce complexity of the parameter space (such as constraining the autoregressive matrices to be identical in all regimes<sup>3</sup>), on the other hand, make the estimation easier, and reliable estimation of such models thereby require less estimation rounds. Constrained estimation is discussed in Section 4.8.

#### 4.7. Estimation of structural STVAR models

As explained, **sstvars** currently supports three types of structural models: structural models identified recursively by the lower triangular Cholesky decomposition, structural models identified by conditional heteroskedasticity, and structural models identified by non-Gaussianity. In either case, the structural models are estimated with the function **fitSSTVAR** based on preliminary estimates from a reduced form model. If the structural model is not overidentifying, which is always the case for recursively identified models, model identified by non-Gaussianity, as well as for models identified by heteroskedasticity when there are two regimes and further constraints are not imposed on the impact matrix, there is no need for estimation but at most for a reparametrization of the model. In any case, the **fitSSTVAR** constructs the structural model appropriately.

To exemplify, we first create a recursively identified structural model based on the reduced form model **fit12** by setting the argument **identification="recursive"**. Then, we will study an example of a structural model identified by heteroskedasticity. The following code builds the recursively identified model:

```
R> fit12rec <- fitSSTVAR(fit12, identification="recursive")
```

Since the parametrization did not change nor was any estimation required, **fit12rec** is essentially the reduced form model **fit12** with has the property that can be used as a structural model in structural analysis such as for estimating the generalized impulse response functions. The following code creates a structural model identified by heteroskedasticity based on the reduced form model **fit12** and then prints it:

```
R> fit12het <- fitSSTVAR(fit12, identification="heteroskedasticity")
R> print(fit12het)
```

logistic Student STVAR model, identified by heteroskedasticity, no AR\_constraints,  
no mean\_constraints, no B\_constraints,

---

<sup>3</sup>Models constrained in this way can often be reliably estimated with a reasonable number of estimation rounds even when  $M > 2$

```
p = 1, M = 2, d = 2, #parameters = 21, #observations = 243 x 2
Switching variable: GDPDEF with lag 1.
```

```
Regime 1
Degrees of freedom: 7.70 (for all regimes)
Regime means: 0.71, 0.49
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 0.63 ] + [ 0.35 -0.35 ] y1.1 + [ 0.37 0.00 ]      eps1
2 y2   [ 0.14 ]   [ 0.06 0.62 ] y2.1   [ 0.00 0.03 ]      eps2
```

```
Regime 2
Weight params: 1.22 (location), 5.01 (scale)
Regime means: 0.77, 1.76
```

```
      Y      phi0      A1      Omega      1/2
1 y1 = [ 2.41 ] + [ 0.13 -0.99 ] y1.1 + [ 1.29 -0.06 ]      eps1
2 y2   [ 0.67 ]   [-0.04 0.64 ] y2.1   [-0.06 0.19 ]      eps2
```

```
Structural parameters:
```

```
      W      lamb2
1 [ 0.17 0.59 ] [ 5.67 ]
2 [ -0.18 0.06 ] , [ 3.29 ]
```

The impact matrix is subject to 0 zero constraints and 0 sign constraints.

Since the structural model identified by heteroskedasticity is not overidentified, no estimation was performed but merely a reparametrization. The estimates for the structural parameters  $W$  and  $\lambda_2, \dots, \lambda_M$  are presented at the bottom of the printout.

If the structural model is identified by heteroskedasticity or non-Gaussianity, additional restrictions can be imposed on the impact matrix by setting them in the argument `B_constraints`. A structural model identified by heteroskedasticity is overidentifying also when there are more than two regimes in the model, as then the matrix decomposition employed in the identification does not always exist. In either case, the structural model needs be estimated, which is performed in `sstvars` based on preliminary estimates obtained either from a reduced form model or from a structural model. The estimation is performed with the function `fitSSTVAR`, which implements a two-phase estimation procedure in which a robust estimation method is used in the first phase and a variable algorithm in the second phase. The default option for the robust method is Nelder-Mead algorithm implemented by [R Core Team \(2022\)](#) in the function `optim` of the package `stats`.

It is important to note that if the initial estimates are far from the ML estimate of the overidentified model, the resulting solution is likely local only due to the high multimodality of the log-likelihood function. However, it is not often very appealing to impose overidentified constraints that far from the unrestricted estimates in the first place. But in any case, since the estimation may be unreliable if the restricted ML estimate is far from the unrestricted ML estimate, we recommend using our package to estimate only such overidentifying structural models in which the unrestricted estimate is close to satisfying the imposed constraints.

The exemplify, we estimate a structural model identified by heteroskedasticity based on the structural model `fit12het` by setting the argument `identification="heteroskedasticity"` and imposing the constraint that the second element of the second column of the impact matrix is zero (the above estimates of  $W$  show that the corresponding unrestricted estimate is close

to zero). The zero constraint is imposed by setting the argument `B_constraints` as matrix such that the second element of the second column is zero and all other elements are `NA`. Sign constraints can be set similarly by setting the corresponding elements to 1 or -1 (or any other strictly positive or negative value). The following code estimates the model:

```
R> fit12hetb <- fitSSTVAR(fit12, identification="heteroskedasticity",
+   B_constraints=matrix(c(NA, NA, NA, 0), nrow=2))
R> print(fit12het)
```

```
The log-likelihood of the supplied model:    -250.236
Constrained log-likelihood prior estimation: -260.164
The log-likelihood after robust estimation:   -250.556
The log-likelihood after final estimation:    -250.486
```

The log-likelihoods of the original model, the initial estimates of the constrained model, the estimates after the robust estimation, and the final estimates are printed. If the log-likelihood of after final estimation is bad compared to the log-likelihood of the original model, the estimation is likely unreliable. The command `print(fit12hetb)` prints the estimated overidentified model (we omit the printout for brevity).

After creating a structural model identified by heteroskedasticity, the columns of  $W$  can be reordered with the function `reorder_B_columns` which also reorders all  $\lambda_{mi}$  accordingly (and hence the resulting model will coincide with the original reduced form model). Also, all signs of any column of  $W$  can be swapped with the function `swap_B_signs`, which also swaps the signs of the corresponding skewness parameter values when `cond_dist = "ind_skewed_t"`. so that the resulting model is observationally equivalent with the original one.

If `cond_dist = "ind_Student"` or `"ind_skewed_t"`, i.e., the shocks are identified by non-Gaussianity, `fitSSTVAR` also enables to conveniently estimate models based on different orderings of signs changes of the columns any of  $B_1, \dots, B_M$ . Such investigation is of interest when the identification is suspected to be weak with respect to the ordering or signs of some of  $B_2, \dots, B_M$  (Virolainen 2025a, see Section 3.3, and for further discussion). Structural models based on different orderings or signs of the columns of  $B_2, \dots, B_M$  can be by specifying the arguments `B_pm_reg`, `B_perm`, and `B_signs` in `fitSSTVAR` accordingly.

#### 4.8. Constrained estimation

`sstvars` supports constrained ML estimation of the STVAR models, including several types of constraints. Linear constraints can be imposed on the autoregressive matrices (argument `AR_constraints`), unconditional means of the regimes can be constrained equal across (groups of) regimes (argument `mean_constraints`), and weight function parameters can be constrained to a fixed value or linear constraints can be impose on them (argument `weight_constraints`). Following sections give examples of constrained estimation imposing some of these constraints.

##### *Linear constraints on the autoregressive parameters*

Imposing linear constraints on the autoregressive parameters of a STVAR model is straightforward in `sstvars`. The constraints are expressed in a somewhat general form which allows to impose a wide class of constraints but one needs to take the time to construct the constraint matrix carefully for each particular case.

We consider constraints of form

$$(\phi_1, \dots, \phi_M) = C\psi, \quad (35)$$

$$\phi_m = (\text{vec}(A_{m,1}), \dots, \text{vec}(A_{m,p})) (pd^2 x_1), \quad m = 1, \dots, M, \quad (36)$$



where  $C$  is known ( $Mpd^2xq$ ) constraint matrix (of full column rank) and  $\psi$  is unknown ( $qx1$ ) parameter vector.

To give couple examples, consider the following two common uses of linear constraints: restricting the autoregressive matrices to be the equal across all regimes and constraining some of the AR parameters to zero.

### Restricting AR matrices to be the equal across the regimes

To restrict the AR matrices to be equal across the regimes, we want  $\phi_m$  to be the same for all  $m = 1, \dots, M$ . The parameter vector  $\psi$  ( $qx1$ ) then corresponds to any  $\phi_m = \phi$ , and therefore  $q = pd^2$ . For the constraint matrix we choose

$$C = [I_{pd^2} : \dots : I_{pd^2}]' \quad (Mpd^2xpd^2), \quad (37)$$

that is,  $M$  pieces of ( $pd^2xpd^2$ ) diagonal matrices stacked on top of each other, because then

$$C\psi = (\psi, \dots, \psi) = (\phi, \dots, \phi). \quad (38)$$

For instance, if there are two regimes in the model, the appropriate constraint matrix then created as

```
R> p <- 1 # The autoregressive order of the model
R> d <- 2 # Whatever the dimension of the time series is
R> I_pd2 <- diag(p*d^2) # The appropriate diagonal matrix
R> (C1 <- rbind(I_pd2, I_pd2)) # Stack them on top of each other
```

	[,1]	[,2]	[,3]	[,4]
[1,]	1	0	0	0
[2,]	0	1	0	0
[3,]	0	0	1	0
[4,]	0	0	0	1
[5,]	1	0	0	0
[6,]	0	1	0	0
[7,]	0	0	1	0
[8,]	0	0	0	1

The command

```
R> fit12c1 <- fitSTVAR(gdpdef, p=1, M=2, weight_function="logistic",
+   weightfun_pars=c(2, 1), cond_dist="Student", AR_constraints=C1,
+   estim_method="two-phase")
```

would then estimate a logistic Student's  $t$  STVAR(1,2) model with first lag of the second variable as the switching variable such that the AR matrices constrained to be the equal in both regimes. We omit the output for brevity. In practice, you might want to adjust the number of CPU cores used (`ncores`), the of estimation rounds (`nrounds`), and set seeds (`seeds`).

### Restricting AR parameters to be the same for all regimes and constraining non-diagonal elements of coefficient matrices to be zero

The previous example shows how to restrict the AR parameters to be the same for all regimes, but say we also want to constrain the non-diagonal elements of coefficient matrices  $A_{m,i}$  ( $m = 1, \dots, M, i = 1, \dots, p$ ) to be zero. We have the constrained parameter  $\psi$  ( $qx1$ ) representing the unconstrained parameters  $(\phi_1, \dots, \phi_M)$ , where the restrictions imply  $\phi_m = \phi = (vec(A_1), \dots, vec(A_p))$  ( $pd^2x1$ ) and the elements of  $vec(A_i)$  ( $i = 1, \dots, p$ ) corresponding to the diagonal are zero.

For illustrative purposes, let's consider a STVAR model with autoregressive degree  $p = 2$ , number of regimes  $M = 2$ , and number of time series in the system  $d = 2$ . Then we have

$$\phi = (A_1(1, 1), 0, 0, A_1(2, 2), A_2(1, 1), 0, 0, A_2(2, 2)) \quad (8 \times 1) \quad \text{and} \quad (39)$$

$$\psi = (A_1(1, 1), A_1(2, 2), A_2(1, 1), A_2(2, 2)) \quad (4 \times 1), \quad (40)$$

where  $A_l(i, j)$  is the  $ij$ th elements of  $A_l$ . By a direct calculation, we can see that choosing the constraint matrix

$$C = \begin{bmatrix} \tilde{c} \\ \tilde{c} \end{bmatrix} \quad (Mpd^2 \times 4), \quad (41)$$

where

$$\tilde{c} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (pd^2 \times 4) \quad (42)$$

satisfies  $C\psi = (\phi, \dots, \phi)$ .

The above constraint matrix can be created as

```
R> c_tilde <- matrix(0, nrow=2*2^2, ncol=4)
R> c_tilde[c(1, 12, 21, 32)] <- 1
R> C2 <- rbind(c_tilde, c_tilde)
R> C2
```

```
      [,1] [,2] [,3] [,4]
[1,]    1    0    0    0
[2,]    0    0    0    0
[3,]    0    0    0    0
[4,]    0    1    0    0
[5,]    0    0    1    0
[6,]    0    0    0    0
[7,]    0    0    0    0
[8,]    0    0    0    1
[9,]    1    0    0    0
[10,]   0    0    0    0
[11,]   0    0    0    0
[12,]   0    1    0    0
[13,]   0    0    1    0
[14,]   0    0    0    0
[15,]   0    0    0    0
[16,]   0    0    0    1
```

The command

```
R> fit12c2 <- fitSTVAR(gdpdef, p=2, M=2, weight_function="logistic",
+   weightfun_pars=c(2, 1), cond_dist="Student", AR_constraints=C2,
+   estim_method="two-phase")
```

would then estimate a logistic Student's  $t$  STVAR(1,2) model with first lag of the second variable as the switching variable such that the AR matrices are constrained to be the equal in both regimes and the off-diagonal elements are restricted to zero. Again, we omit the output for brevity (and you may want to adjust the arguments `nrounds`, `ncores`, and `seeds` when estimating the model in practice).

### *Constraining the unconditional means of some regimes to be equal*

In addition to constraining the autoregressive parameters, `sstvars` allows to constrain the unconditional means of some regimes to be the equal. This feature is, however, only available for models that are parametrized with the unconditional means instead of intercepts (because some of the estimation is always done with mean-parametrization and one cannot generally swap the parametrization when constraints are imposed on means/intercepts). With the mean-parametrization employed (by setting `parametrization="mean"`), one may define groups of regimes that have the same mean parameters using the argument `mean_constraints`. For instance, with three regime model ( $M = 3$ ) the argument `mean_constraints=list(c(1, 3), 2)` sets the unconditional means of the first and third regimes to be the same while allows the second regime to have different mean.

One can also combine linear constraints on the AR parameters with constraining some of the means to be the same. This allows, for instance, to estimate a model in which only the covariance matrix varies in time. To exemplify, the following code estimates a Student's  $t$  logistic STVAR( $p = 1, M = 2$ ) model such that the unconditional means and autoregression matrices are constrained be the same in both regimes. The resulting model thereby has time-varying covariance matrix but otherwise it is linear.

```
R> fit12c3 <- fitSTVAR(gdpdef, p=1, M=2, weight_function="logistic",
+   weightfun_pars=c(2, 1), cond_dist="Student", AR_constraints=C1,
+   mean_constraints=list(1:2), parametrization="mean",
+   estim_method="two-phase")
```

The output is omitted for brevity.

### *Constraining weight functions parameters*

It is also possible to constrain the weight functions parameters  $\alpha$  (e.g., the location and scale parameters for logistic models). `sstvars` accommodates two types of alternative constraints on the weight function parameters: linear constraints and fixed values. Note that weights constraints are not available for models with exogenous weights, as they do not contain any weight function parameters.

#### **Linear constraints**

Linear constraints on the weight function parameters are of the form

$$\alpha = R\xi + r, \quad (43)$$

where  $\alpha$  ( $a \times 1$ ) contains the weight function parameters,  $R$  is a known ( $a \times l$ ) constraint matrix of full column rank,  $r$  is a known ( $a \times 1$ ) constant, and  $\xi$  is an unknown ( $l \times 1$ ) parameter. The constraint matrix  $R$  and the constant  $r$  are set in the argument `weight_constraints` as a list of two elements,  $R$  in the first element and  $r$  in the second element. The number of unknown parameters  $l$  is the number of columns of  $R$ . For instance, the following argument imposes constraints for the scale and location parameters of a logistic STVAR model (in which  $\alpha = (c, \gamma)$ ) such that the location parameter is the scale parameter divided by two plus 0.3: `weight_constraints=list(R=matrix(c(0.5, 1), nrow=2), r=c(0.3, 0))`.

### Fixed values

Imposing the weight function parameters to be known fixed values is very straightforward. In this case, the argument `weight_constraints` should still be a list including the elements  $R$  and  $r$ , but the former is set to zero,  $R = 0$ , and the latter is set to the desired fixed values. For instance the following argument imposes constraints for the location and location parameters of a logistic STVAR model (in which  $\alpha = (c, \gamma)$ ) such that the location parameter is 0.3 and the scale parameter is 0.5: `weight_constraints=list(R=0, r=c(0.3, 0.5))`.

## 5. Testing parameter constraints

One way to assess the validity of the imposed constraints is to compare the values of information criteria of the constrained and unconstrained models. `sstvars`, however, also provides functions for testing the constraints with the likelihood ratio test, Wald test, and Rao's test, whose applicability requires that the ML estimator of the STVAR model has the conventional asymptotic distribution. As noted before, this is a mere assumption, but given the process is ergodic stationary, there is no particular reason to believe that the standard asymptotic results would not hold. For a discussion on the tests, see Buse (1982) and the references therein, for example.

The likelihood ratio test considers the null hypothesis that the true parameter value  $\theta_0$  satisfies some constraints imposed on these parameters (such that the constrained parameter space is a subset of the parameter space, which is presented in Assumption 2). Denoting by  $\hat{L}_U$  and  $\hat{L}_C$  the (maximized) log-likelihoods based on the unconstrained and constrained ML estimates, respectively, the test statistic takes the form

$$LR = 2(\hat{L}_U - \hat{L}_C). \quad (44)$$

Under the null, the test statistic is asymptotically  $\chi^2$ -distributed with the degrees of freedom given by the difference in the dimensions of the unconstrained and constrained parameter spaces. With `sstvars`, the likelihood ratio test can be calculated with the function `LR_test`, which takes the unconstrained model (a class '`stvar`' object) as its first argument and the constrained model as the second argument.

`sstvars` implements the Wald test of the null hypothesis

$$A\theta_0 = c, \quad (45)$$

where  $A$  is a  $(k \times d)$  matrix with full row rank,  $c$  is a  $(k \times 1)$  vector,  $\theta_0$  is the true parameter value,  $d$  is the dimension of the parameter space, and  $k$  is the number of constraints. The Wald test statistic takes the form

$$W = (A\hat{\theta} - c)'[A\mathcal{J}(\hat{\theta})^{-1}A']^{-1}(A\hat{\theta} - c), \quad (46)$$

where  $\mathcal{J}(\hat{\theta})$  is the observed information matrix evaluated at the ML estimate  $\hat{\theta}$ . Under the null, the test statistic is asymptotically  $\chi^2$ -distributed with  $k$  degrees of freedom (which is the difference in dimensions of the constrained and unconstrained parameter spaces). With `sstvars`, the Wald test can be calculated with function `Wald_test`, which takes the estimated unconstrained model (as a class '`stvar`' object) as the first argument, the matrix  $A$  as the second argument, and the vector  $c$  as the third argument.

Finally, Rao's test is implemented to the function `Rao_test`. See the function documentation on how to use it.

## 6. Residual based model diagnostics

**sstvars** employs residual based diagnostics for assessing the adequacy of the fitted model. Conventional graphical diagnostics can be examined with function `diagnostic_plot`, which plots the residual time series, auto- and crosscorrelation functions of the residuals, auto- and cross-correlation functions of the squared residuals, and normal quantile-quantile plots as well as histograms of the residuals. The plots can be created for both unstandardized residuals or for standardized residuals by adjusting the argument `standardize` to `FALSE` or `TRUE` according. Using unstandardized residuals is advisable when checking for remaining autocorrelation. But standardized residuals should be used to check for remaining conditional heteroskedasticity and to check the model's adequacy to capture the marginal distribution of the series, because the STVAR models are conditionally heteroskedastic and the unstandardized residuals do not take into account the time-varying conditional covariance matrix.

Remaining autocorrelation in the residuals can also be tested with the (adjusted) Portmanteau test, which is implemented to the function `Portmantau_test`. The test can also be applied to standardized squared residuals to test for remaining conditional heteroskedasticity by setting the argument `which_test="het.sked"`. The number of lags that should be taken into account in the test is set with the argument `nlags`.

## 7. Impulse response analysis

### 7.1. Generalized impulse response function

The expected effects of the structural shocks in the structural STVAR models generally depend on the initial values as well as on the sign and size of the shock, which makes the conventional way of calculating impulse responses unsuitable (see, e.g., Kilian and Lütkepohl 2017, Chapter 4). Therefore, we consider the generalized impulse response function (GIRF) (Koop *et al.* 1996) defined as

$$\text{GIRF}(n, \delta_j, \mathcal{F}_{t-1}) = E[y_{t+n} | \delta_j, \mathcal{F}_{t-1}] - E[y_{t+n} | \mathcal{F}_{t-1}], \quad (47)$$

where  $n$  is the chosen horizon,  $\mathcal{F}_{t-1} = \sigma\{y_{t-j}, j > 0\}$  as before, the first term on the right side is the expected realization of the process at time  $t + n$  conditionally on a structural shock of sign and size  $\delta_j \in \mathbb{R}$  in the  $j$ th element of  $e_t$  at time  $t$  and the previous observations, and the second term on the right side is the expected realization of the process conditionally on the previous observations only. GIRF thus expresses the expected difference in the future outcomes when the specific structural shock hits the system at time  $t$  as opposed to all shocks being random.

Due to the  $p$ -step Markov property of the implemented STVAR models, conditioning on (the  $\sigma$ -algebra generated by) the  $p$  previous observations  $\mathbf{y}_{t-1} \equiv (y_{t-1}, \dots, y_{t-p})$  is effectively the same as conditioning on  $\mathcal{F}_{t-1}$  at the time  $t$  and later. The initial values (or history)  $\mathbf{y}_{t-1}$  can be either fixed or random, but with random history the GIRF becomes a random vector, however. Using fixed  $\mathbf{y}_{t-1}$  makes sense when one is interested in the effects of the shock in a particular point of time. Alternatively, one can estimate GIRFs conditional on the initial values being from a specific regime, in which case  $\mathbf{y}_{t-1}$  should be generated from the regime of interest.

In practice, the GIRF and its distributional properties can be approximated with a Monte Carlo algorithm that generates independent realizations of the process and then takes the sample mean for point estimate. If  $\mathbf{y}_{t-1}$  is random and follows the distribution  $G$ , the GIRF should be estimated for different values of  $\mathbf{y}_{t-1}$  generated from  $G$ , and then the sample mean and sample quantiles can be taken to obtain the point estimate and confidence intervals. **sstvars** accommodates generating the initial values from the full STVAR process or from a given regime. Alternatively, the GIRF can be estimated for a set of fixed initial values, such as for the length  $p$  histories in the data. This can be done by setting the argument `use_data_shocks=TRUE`, in which case, the related signs and sizes of the structural shocks recovered from the data are also

in calculation of the GIRF, thereby reflecting the properties of the data (adjust the argument `data_girf_pars` to adjust whether a specific regime should be dominant in the histories, or whether only shocks of specific sign or size should be used). The algorithm for calculating the GIRFs implemented in `sstvars` is presented in [Lanne and Virolainen \(2025\)](#).

Because the STVAR models allow to associate specific features or economic interpretations for different regimes, and because shifts in the regime are the source of asymmetries in the impulse responses, it might be interesting to also examine the effects of a structural shock to the transition weights  $\alpha_{m,t}$ ,  $m = 1, \dots, M$ . We then consider the related GIRFs  $E[\alpha_{m,t+n}|\delta_j, \mathbf{y}_{t-1}] - E[\alpha_{m,t+n}|\mathbf{y}_{t-1}]$  for which point estimates and confidence intervals can be constructed similarly to (47).

In `sstvars`, the GIRF can be estimated with the function `GIRF` which should be supplied with the estimated STVAR model or a STVAR model built with hand-specified parameter values using the function `STVAR`. Structural models can be created based on a reduced form model with the function `fitSSTVAR`. The sign and size of the structural shock can be set with the argument `shock_size` (ignored if `use_data_shocks=TRUE`). If not specified, a positive shock with the size of one standard deviation is used; that is, the size is one. Among other arguments, the function may also be supplied with the argument `init_regime` which specifies from which regime the initial values are generated from (ignored if `use_data_shocks=TRUE`). Alternatively, one may specify fixed initial values with the argument `init_values` (ignored if `use_data_shocks=TRUE`), or use initial values in the observed data with `use_data_shocks=TRUE` as described above. Note that the confidence intervals (whose confidence level can be specified with the argument `ci`) reflect uncertainty about the initial value only and not about the parameter estimates.

Due to the nonlinear nature of the model, GIRFs estimated from different starting values, or with different sign or magnitude of the shock, generally move the variables differently. Sometimes it is, however, of interest to scale the impulse responses so that they correspond to instantaneous movement of some specific sign and size of some specific variable. In `sstvars`, this is most conveniently achieved with the arguments `scale`. The argument `scale` can be specified in order to scale the GIRFs to some of the shocks so that they correspond to a specific sign and size of instantaneous response of some specific variable. Alternatively, the GIRFs can be scaled to correspond to a specific peak response of some variable by setting `scale_type="peak"`. For a single shock, the argument `scale` should be a length three vector where the shock of interest is given in the first element (an integer in  $1, \dots, d$ ), the variable according to which the GIRFs should be scaled in the second element (an integer in  $1, \dots, d$ ), and the sign and size of the given variable's instantaneous response in the third element (a non-zero real number). If the GIRFs of multiple shocks should be scaled, provide a matrix which has one column for each of the shocks with the columns being the length three vectors described above. Note that if you scale the GIRFs, the scaled GIRFs of transition weights can be outside the interval from zero to one.

Because estimating the GIRF and their confidence intervals is computationally demanding, parallel computing is taken use of to shorten the estimation time. The number of CPU cores used can be set with the argument `ncores`. The objects returned by the `GIRF` function have their own `plot` and `print` methods. Also, cumulative impulse responses of the specified variables can be obtained directly by specifying the argument `which_cumulative`.

## 7.2. Generalized forecast error variance decomposition

Similarly to the conventional impulse response functions are unsuitable for impulse response analysis (due to their inability to capture asymmetries the effects of the shocks), the conventional forecast error variance decomposition is unsuitable for tracking the contribution of each shock to the variance of the forecast errors. We consider the generalized forecast error variance decomposition (GFEVD) ([Lanne and Nyberg 2016](#)) that is defined for variable  $i$ , shock  $j$ , and

horizon  $h$  as

$$\text{GFEVD}(j, y_{it}, \delta_j, \mathcal{F}_{t-1}) = \frac{\sum_{l=0}^h \text{GIRF}(l, \delta_j, \mathcal{F}_{t-1})_i^2}{\sum_{k=1}^d \sum_{l=0}^h \text{GIRF}(l, \delta_k, \mathcal{F}_{t-1})_i^2}, \quad (48)$$

where  $h$  is the chosen horizon and  $\text{GIRF}(l, \delta_j, \mathcal{F}_{t-1})_i$  is the  $i$ th element of the related GIRF (see also the notation described for GIRF in the previous section). That is, the GFEVD is otherwise similar to the conventional forecast error variance decomposition but with GIRFs in the place of conventional impulse response functions. Because the GFEVDs sum to unity (for each variable), they can be interpreted in a similar manner to the conventional FEVD.

In **sstvars**, the GFEVD can be estimated with the function **GFEVD**. As with the GIRF, the GFEVD is dependent on the initial values. The type of the initial values is set with the argument **initval\_type**, and there are three options:

1. **"data"** which estimates the GFEVDs for all, or a subset of, possible length  $p$  histories in the data, and then the final GFEVD is obtained as the sample mean over them.
2. **"random"** which generates the initial values from one of the specific regimes, specified by the argument **init\_regimes**. The GFEVD is calculated for each initial value, and then the final GFEVD is obtained as the sample mean over them.
3. **"fixed"** which estimates the GFEVD for a single fixed initial value that is set with the argument **init\_values**.

The shock size is the same for all scalar components of the structural shock and it can be adjusted with the argument **shock\_size**. If the GIRFs for some variables should be cumulative before calculating the GFEVD, specify them with the argument **which\_cumulative**. Finally, note that the GFEVD objects have their own plot and print methods.

**sstvars** also implements a special feature in which for every possible length  $p$  history in the data, the GFEVD is estimated for a shock that has the sign and size of the corresponding structural shock recovered from the fitted model. This can be done by setting the argument **use\_data\_shocks=TRUE**. The GFEVD is then calculated as the average of the GFEVDs obtained from the GIRFs estimated for the data shocks. The plot and print methods can be used as usual for this GFEVD. However, this feature also estimates the contribution of each shock to the variance of the forecast errors at various horizons in specific historical points of time. This can be done by using the plot method with the argument **data\_shock\_pars**. Note that the arguments **shock\_size**, **initval\_type**, and **init\_regime** are ignored if **use\_data\_shocks == TRUE**. Use the argument **data\_gfevd\_pars** to further adjust the settings with **initval\_type="data"** or **use\_data\_shocks=TRUE**.

### 7.3. Linear impulse response functions

It is also possible to calculate linear impulse response functions (IRF) based on a specific regime of the estimated model by using the function **linear\_IRF**. If the autoregressive dynamics of the model are linear (i.e., either  $M = 1$  or mean and AR parameters are constrained identical across the regimes), confidence bounds can be estimated based on a type of a fixed-design wild residual bootstrap method. **sstvars** implements the method proposed [Herwartz and Lütkepohl \(2014\)](#).

## 8. Historical decompositions

**sstvars** implements the historical decomposition of the observed time series into the contributions of the structural shocks, initial conditions, and so-called "steady state component" as



described in [Wong \(2018\)](#). To obtain the historical decomposition, it is convenient to reformulate the STVAR model in the VAR(1) companion form. First, note that the model (1) can be written as

$$y_t = \phi_{y,t} + \sum_{i=1}^p A_{y,t,i} y_{t-i} + B_{y,t} e_t, \quad (49)$$

where  $\phi_{y,t} \equiv \sum_{m=1}^M \alpha_{m,t} \phi_m$  ( $d \times 1$ ) and  $A_{y,t,i} \equiv \sum_{m=1}^M \alpha_{m,t} A_{m,i}$  contain the time-varying AR coefficients. Depending on the shock distributions and the identification method, the impact matrix  $B_{y,t}$  is either directly parametrized to the model or obtained at each  $t$  by decomposing the conditional covariance matrix  $\Omega_{y,t}$  accordingly.

Denoting  $\mathbf{y}_t \equiv (y_t, \dots, y_{t-p+1})$  ( $pd \times 1$ ), the model can then be written in the companion form as

$$\mathbf{y}_t = H\phi_{y,t} + A_{y,t}\mathbf{y}_{t-1} + HB_{y,t}e_t, \quad (50)$$

where

$$H = \begin{bmatrix} I_d \\ \mathbf{0}_{d(p-1) \times d} \end{bmatrix}_{(pd \times d)}, \quad A_{y,t} = \begin{bmatrix} A_{y,t,1} & \cdots & A_{y,t,p} \\ I_{d(p-1)} & & \mathbf{0}_{d(p-1) \times d} \end{bmatrix}_{(pd \times pd)}, \quad (51)$$

$I_d$  is the ( $d \times d$ ) diagonal matrix, and  $\mathbf{0}_{d(p-1) \times d}$  is the  $(d(p-1) \times d)$  dimensional matrix of zeros. Suppose the observed time series is indexed as  $y_{t-p+1}, \dots, y_0, y_1, \dots, y_T$ . As described in [Wong \(2018, Section 2\)](#), recursive substitution of Equation (50) yields:

$$\begin{aligned} \mathbf{y}_t = & \underbrace{HB_{y,t}e_t + \sum_{j=1}^{t-1} \left( \prod_{i=0}^{j-1} A_{y,t-i} \right) HB_{y,t}e_{t-j}}_{\text{Contribution of the shocks}} + \\ & \underbrace{\left( \prod_{i=0}^{t-1} A_{y,t-i} \right) \mathbf{y}_0}_{\text{Initial conditions}} + \underbrace{H\phi_{y,t} + \sum_{j=1}^{t-1} \left( \prod_{i=0}^{j-1} A_{y,t-i} \right) H\phi_{y,t-j}}_{\text{Steady state component}}. \end{aligned} \quad (52)$$

Baseline projection

The cumulative contributions of the shocks to the fluctuations of the variables are obtained from the block "Contribution of the shocks" in the above equation, whereas the effect of the initial conditions are obtained from the block "Initial conditions". The block "Steady state component" contains so-called steady-state component of the historical decomposition, which is discussed in [Wong \(2018\)](#), Section 3.2.

In `sstvars`, the historical decomposition is computed with the function `hist_decomp`, which should be supplied by a fitted STVAR model (a class `stvar` object). The returned historical decomposition object has its own plot and print methods for convenient examination of the contributions of the shocks. While included in the returned object, the initial condition and steady state -components are not included in the print and plot methods, however.

## 9. Counterfactual analysis

Counterfactual analysis addresses questions of the form "What would (have) happen(ed) if...?", and many different types of counterfactuals have been discussed in the literature (see, e.g., [Kilian and Lütkepohl 2017](#), Chapter 4, and the references therein). Two different types of policy counterfactuals are implemented to `sstvars`: counterfactuals in which the policy variable takes some given path in given points of given (e.g., [Bernanke, Gertler, and Watson 1997](#)) and counterfactuals in which the policy variable does not response to lagged movements of some

given variable nor to shock to it (or either) in given points of time (e.g., Kilian and Lewis 2011). Following Bernanke *et al.* (1997), Kilian and Lewis (2011), and others, we simulate the counterfactual scenarios by creating hypothetical shocks to the policy variable that yield the counterfactual outcome. This approach has the appealing feature that counterfactual deviations from the policy reaction function are treated as policy surprises, allowing them to propagate normally. Thus, the dynamics of the model are not, *per se*, tampered but just the policy surprises are.

Consider first counterfactual scenarios in which the policy variable of interest takes some given values in a given points of time. Here, it is shown for a single time period how the shock to policy variable that sets it take a given value is found. In the following sections, the algorithms simulating historical counterfactuals, counterfactual forecast scenarios, and counterfactual GIRFs are established by making use the single time period case iteratively.

Consider the counterfactual scenario in which the  $i_1$ th variable takes some specific value in the time period  $t$ , and denote this counterfactual value of the  $i_1$ th variable as  $y_{i_1t}^* \in \mathbb{R}$ . To simulate the counterfactual scenario in which this observation of the  $i_1$ th variable is achieved, we follow Bernanke *et al.* (1997) and create a hypothetical shock to it that sets  $y_{i_1t} = y_{i_1t}^*$ . Assuming that the  $i_1$ th shock is the shock the  $i_1$ th variable, the task is then to manipulate the value of  $e_{i_1t}$  to obtained the desired observation of the  $i_1$ th variable (the policy variable of interest). For recursively identified models, this should be automatically satisfied. In models identified by heteroskedasticity or non-Gaussianity, the ordering of the shocks can generally be changed without loss of generality, in which case the ordering be changed with the function `reorder_B_columns` so that the  $i_1$ th shock is the shock to the  $i_1$ th variable.

Consider the STVAR model (1) in the form (49). The (estimated) values of  $\phi_{y,t}$ ,  $A_{y,t,1}, \dots, A_{y,t,p}$ , and  $B_{y,t}$  are known by assumption conditionally on  $\mathcal{F}_{t-1}$ . Therefore, for a given time period  $t$ , the value of  $e_{i_1t}$  that sets  $y_{i_1t} = y_{i_1t}^*$  is found as a solution to the single linear equation obtained from the  $i$ th row of (49) with  $y_{i_1t} = y_{i_1t}^*$  substituted:

$$y_{i_1t}^* = \mu_{t,i_1} + \sum_{j=1}^d b_{t,i_1j} e_{jt}, \quad (53)$$

where  $\mu_{t,i_1}$  is the  $i_1$ th element of the vector  $\phi_{y,t} + \sum_{i=1}^p A_{y,t,i} y_{t-i}$  and  $b_{t,i_1j}$  is the  $i_1j$ th element of the impact matrix  $B_{y,t}$ . Solving this equation with respect to  $e_{i_1t}$  gives

$$e_{i_1t} = b_{t,i_1i_1}^{-1} \left( y_{i_1t}^* - \mu_{t,i_1} - \sum_{j=1, j \neq i_1}^d b_{t,i_1j} e_{jt} \right), \quad (54)$$

where it is assumed that  $b_{t,i_1i_1} \neq 0$  (i.e., that the shock to the policy variable affects the policy on impact). Denote this solution of  $e_{i_1t}$  as  $e_{i_1t}^*$ . The implied counterfactual time  $t$  observations are obtained by replacing the  $i_1$ th shock in the vector  $e_t$  by  $e_{i_1t}^*$ , and then calculating the implied observations from (49) by using the obtained hypothetical structural error.

Consider then a counterfactual scenario in which the response of the policy variable is shut off to lagged and contemporaneous movements of a given variable, but policy variable is allowed to respond normally to movements of the other included included variables. Following Kilian and Lewis (2011), we the simulate the counterfactual scenario by creating hypothetical shocks to the policy variable that off set the response of the policy variable to the movements of a given variable. Suppose the policy variable of interest is the  $i_1$ th variable and variable whose movements the policy variable should not react to at the time  $t$  is the  $i_2$ th variable satisfying  $i_2 \neq i_1$ . Assume that the  $i_1$ th shock is the shock to the  $i_1$ th variable and that the  $i_2$ th shock is the shock to the  $i_2$ th variable (again, this should be automatically satisfied for recursively identified models, whereas the function `reorder_B_columns` can be used to change the ordering

of the shocks for model identified by heteroskedasticity or non-Gaussianity).<sup>4</sup>

The lagged and contemporaneous effects of the  $i_2$ th variable to the  $i_1$ th variable are

$$\sum_{i=1}^p a_{t,i,i_1i_2} y_{i_2t-i} + b_{t,i_1i_2} e_{i_2t} \quad (55)$$

where  $a_{t,i,i_1i_2}$  is the  $i_1i_2$ th element of the coefficient matrix  $A_{t,i}$  and  $b_{t,i_1i_2}$  is the  $i_1i_2$ th element of the impact matrix  $B_t$ , (estimated) values of which are, conditionally on  $\mathcal{F}_{t-1}$ , by assumption known at the time  $t$ . Thus, subtracting the quantity

$$b_{t,i_1i_1}^{-1} \left( \sum_{i=1}^p a_{t,i,i_1i_2} y_{i_2t-i} + b_{t,i_1i_2} e_{i_2t} \right) \quad (56)$$

from the  $i_1$ th shock  $e_{i_1t}$  shuts off these effects from the movements of the  $i_1$ th variable (assuming  $b_{t,i_1i_1} \neq 0$ ).

Now, to simulate the counterfactual scenario in which the response of the  $i_1$ th variable does not respond to the lagged and contemporaneous movements of the  $i_2$ th variable, is obtained by first replacing the  $i_1$ th shock in the vector  $e_t$  by

$$e_{i_1t}^* = e_{i_1t} - b_{t,i_1i_1}^{-1} \left( \sum_{i=1}^p a_{t,i,i_1i_2} y_{i_2t-i} + b_{t,i_1i_2} e_{i_2t} \right), \quad (57)$$

where  $e_{i_1t}$  is the  $i_1$ th element of  $e_t$  before manipulating it. Then, the implied counterfactual observations are calculated from (49) by using the obtained hypothetical structural error.

Next, we show how the above techniques derived for a single time period can be applied iteratively to simulate several types of counterfactual scenarios, including historical counterfactuals, forecast scenarios, and counterfactual GIRFs.

### 9.1. Historical counterfactuals

We consider historical policy counterfactual scenarios where, in given historical points of time, the policy variable(s) of interest takes some hypothetical path, or its responses to lagged and contemporaneous movements of some given variable is shut off. Suppose the observed time series is  $y_{t-p+1}, \dots, y_0, y_1, \dots, y_T$ . Consider a counterfactual scenario in which the  $i_1$ th variable either takes some specific values, or its responses to lagged and contemporaneous movements of the  $i_2$ th variable ( $i_2 \neq i_1$ ) are shut off, in the time periods  $t^*, t^*+1, \dots, t^*+k$ ,  $t^* \in \{1, \dots, T-k\}$ ,  $k \leq T - t^*$ . Assuming that the  $i_1$ th shock is the shock the  $i_1$ th variable, the task is then to manipulate the values of  $e_{i_1t^*}, \dots, e_{i_1t^*+k}$ , with the rest of the shocks are taken to be the structural shocks recovered from the (fitted) model, to obtain the desired counterfactual scenario. In the latter scenario we also assume that the  $i_2$ th shock is the shock to the  $i_2$ th variable.

Using the results derived in Section 9, the counterfactual scenario can be simulated iteratively with the following algorithm:

The resulting counterfactual path of the observable variables represents the model implied scenario in which, *ceteris paribus*, the policy variable either takes the specified path or does not respond to movements of the  $i_2$ th variable (depending on the considered scenario) in the time periods  $t^*, \dots, t^*+k$ . In **sstvars**, the historical counterfactuals can be simulated with the function `cfact_hist`.

### 9.2. Forecast scenarios

---

<sup>4</sup>If it is not clear whether the  $i_2$ th shock of interest can be interpreted as a shock to a variable (but has a broader definition such as "a demand shock"), the counterfactual scenario is interpreted as follows: the  $i_1$ th variable does not react to lagged movements of the  $i_2$ th variable nor to the  $i_2$ th shock.

**Algorithm 1** Historical counterfactual

- 
- 1: **for** each  $t$  in  $t^*, \dots, T$  **do**
  - 2:   Compute the values of  $\phi_t$ ,  $A_{t,1}, \dots, A_{t,p}$ , and  $B_t$  based on the preceding observations, and take the structural error  $e_t$  recovered from the (fitted) model.
  - 3:   **if**  $t \in \{t^*, \dots, t^* + k\}$  **then**
  - 4:     Calculate the counterfactual shock  $e_{i_1 t}^*$  from (54) or (56) depending on the type of the counterfactual, and replace the  $i_1$ th element of  $e_t$  by this hypothetical shock.
  - 5:   **end if**
  - 6:   Take the structural error  $e_t$  obtained from the previous steps and compute the corresponding (counterfactual) observation  $y_t$  from (49).
  - 7: **end for**
- 

Forecast scenarios refer to alternative predictions of future outcomes based on different hypothetical conditions or assumptions. For instance, when evaluating economic policy, one might create a baseline forecast assuming no policy change, and alternative scenarios assuming specific policy implementations. We consider the type of forecast scenarios in which a policy variable either takes some specific values at given forecast horizons (as opposed to its predicted values without such assumptions) or its responses to lagged and contemporaneous movements of some given variable are shut off.

Deriving multistep predictions and prediction intervals analytically for nonlinear SVAR models can be challenging, as the dynamics of the model may vary depending the realized observations. Therefore, we consider simulation based forecasting in which a large number of potential future sample paths of the process is simulated, and the sample mean (or median) over the Monte Carlo (MC) repetitions is taken as the point prediction and sample quantiles as the prediction intervals. See Section 11 discussion about forecasting by simulation. The counterfactual forecast scenario is created by manipulating the shocks to the policy variable in each MC repetition.

To formally describe the method, consider the counterfactual scenario in which the  $i_1$ th variable either takes some specific values, or its responses to the movements the  $i_2$ th variable ( $i_2 \neq i_1$ ) are shut off, in the forecast horizons  $h^*, h^* + 1, \dots, h^* + k$ , where  $h^* > 0$  and  $k \geq 0$ . Again, we assume that the  $i_1$ th shock is the shock to the  $i_1$ th variable (and in the latter type of scenario also that the  $i_2$ th shock is the shock to the  $i_2$ th variable). Suppose that the forecast is based on the information up to the time  $T$ . The forecast scenario is simulated iteratively with the following steps:

**Algorithm 2** Counterfactual forecast scenario

- 
- 1: Set the horizon  $H$  and the number of repetitions  $R$ .
  - 2: **for** each  $h$  in  $1, 2, \dots, H$  **do**
  - 3:   Conditionally on  $\mathcal{F}_{T+h-1}$ , draw  $e_{T+h}$  from its (estimated) distribution.
  - 4:   **if**  $h \in \{h^*, \dots, h^* + k\}$  **then**
  - 5:     Calculate the counterfactual shock  $e_{i_1 T+h}^*$  from (54) or (56) depending on the type of the counterfactual, and replace the  $i_1$ th element of  $e_{T+h}$  by this hypothetical shock.
  - 6:   **end if**
  - 7:   Conditionally on  $\mathcal{F}_{T+h-1}$  and  $e_{T+h}$  obtained from the previous steps, compute the corresponding observation  $y_{T+h}$  from (49).
  - 8: **end for**
  - 9: Repeat Steps 2-8  $R$  times, and the sample mean or median over the MC repetitions for the point prediction and desired sample quantiles for the prediction intervals.
- 

The baseline (not counterfactual) point prediction and prediction intervals are obtained from Algorithm 2 by dropping Steps 4-6 from it. See also Section 11. In **sstvars**, the counterfactual

forecast scenarios can be simulated with the function `cfact_fore`.

### 9.3. Counterfactual GIRFs

It can also be interesting to incorporate counterfactual scenarios impulse response analysis, to assess what the effects a shock if the policy variable moves in a some specific way. Similarly to the historical counterfactuals and forecast scenarios, we consider the type of counterfactual GIRFs in which a policy variable either takes some specific values at given GIRF horizons (as opposed to its simulated values without such assumptions) or its responses to lagged and contemporaneous movements of some given variable are shut off. In particular, we simulate the counterfactuals by creating hypothetical shocks to the policy variable that yield the counterfactual outcome. See Section 7.1 for the definition of GIRFs.

Consider a counterfactual scenario in which the  $i_1$ th variable either takes some specific values, or its responses to lagged and contemporaneous movements of the  $i_2$ th variable ( $i_2 \neq 1$ ) is shut off, in the GIRF horizons  $h^*, h^* + 1, \dots, h^* + k$ , where  $h^* \geq 0$  and  $k \geq 0$ . In the following, for the  $l$ th Monte Carlo repetition,  $y_{t+h}^{(l)}(\delta_j, \mathbf{y}_{t-1})$  denotes a realization of the process at time  $t + h$  conditional on the history  $\mathbf{y}_{t-1}$  and the  $j$ th structural shock of sign and size  $\delta_j \in \mathbb{R}$  arriving at time  $t$ , whereas  $y_{t+h}^{(l)}(\mathbf{y}_{t-1})$  denotes an alternative realization conditional on the history  $\mathbf{y}_{t-1}$  only.

For a single history  $\mathbf{y}_{t-1}$  and given sign and size  $\delta_j$  of the  $j$ th structural shock, the counterfactual GIRF can be estimated with the following algorithm (adapted from the GIRF algorithm in Lanne and Virolainen 2025, Appendix B), where we assume that the shock  $j$  is not the shock to the policy variable, i.e.,  $j \neq i_1$ :

The GIRF that does not impose a counterfactual scenario is obtained from Algorithm 3 by simply dropping the Steps 7-12 from it. See also Section 7.1. In `sstvars`, the counterfactual GIRFs can be simulated with the function `cfact_girf`.

## 10. Building a STVAR model with specific parameter values

The function `STVAR` facilitates building STVAR models without estimation, for instance, in order to simulate observations from a STVAR process with specific parameter values. The function should be supplied at least with the arguments `p`, `M`, and `d` specifying the autoregressive order, the number of regimes, and the dimension of the time series, respectively. The argument `params` should be used to specify the parameter values, whereas the weight function and weight function parameters are specified in the arguments `weight_function` and `weightfun_pars`, respectively, and the conditional distribution in the function `cond_dist`.

To exemplify, we build a reduced form Gaussian STVAR  $p = 1$ ,  $M = 1$ ,  $d = 2$  model with relative stationary densities as the transition weights. The model has intercept parametrization and parameter values  $\varphi_{1,0} = (0, 1)$ ,  $\varphi_{12,0} = (0, 2)$ ,  $\text{vec}(A_{1,1}) = (0.2, 0.2, 0.2, -0.2)$ ,  $\text{vec}(A_{1,1}) = (0.3, 0.3, 0.3, -0.3)$ ,  $\text{vech}(\Omega_1) = (1, 0.1, 1)$ ,  $\text{vech}(\Omega_2) = (4, 0.4, 4)$ , and  $\alpha_1 = 0.6$ . After building the model, we use the `print` method to examine it:

```
R> params122 <- c(0, 1, 0, 2, 0.2, 0.2, 0.2, -0.2, 0.3, 0.3, 0.3, -0.3, 1,
+ 0.1, 1, 4, 0.4, 4, 0.6)
R> mod122 <- STVAR(p=1, M=2, d=2, params=params122,
+ weight_function="relative_dens")
R> mod122
```

```
relative_dens Gaussian STVAR model, reduced form model, no AR_constraints,
no mean_constraints,
```

**Algorithm 3** Counterfactual GIRF

- 
- 1: Set the horizon  $H$  and the number of repetitions  $R$ .
  - 2: **for** each  $h$  in  $0, 1, \dots, H$  **do**
  - 3:   Conditionally on  $\mathcal{F}_{t+h-1}$ , draw  $e_{t+h}$  from its (estimated) distribution.
  - 4:   **if**  $h = 0$  **then**
  - 5:     Impose the sign and size  $\delta_j$  to the  $j$ th element of  $e_{t+h}$  to obtain the modified structural error  $\tilde{e}_{t+h}$ .
  - 6:   **end if**
  - 7:   **if**  $h \in \{h^*, \dots, h^* + k\}$  **then**
  - 8:     Calculate the counterfactual shock  $e_{i_1 t+h}^*$  from (54) or (56) depending on the type of the counterfactual, and replace the  $i_1$ th element of  $e_{t+h}$  by this hypothetical shock.
  - 9:     **if**  $h = 0$  **then**
  - 10:      Calculate also the modified counterfactual shock  $\tilde{e}_{i_1 t+h}^*$  using the modified structural error  $\tilde{e}_{t+h}$  in (54) or (56) depending on the type of the counterfactual, and replace the  $i_1$ th element of  $\tilde{e}_{t+h}$  by this hypothetical shock.
  - 11:     **end if**
  - 12:   **end if**
  - 13:   Calculate  $y_{t+h}^{(l)}(\delta_j, \mathbf{y}_{t-1})$  by using the previous observations  $(\mathbf{y}_{t-1}, y_t^{(l)}(\delta_j, \mathbf{y}_{t-1}), \dots, y_{t+h-1}^{(l)}(\delta_j, \mathbf{y}_{t-1})$  and the structural error  $e_{t+h}$  (or  $\tilde{e}_{t+h}$  if  $h = 0$ ) obtained from Steps 3-12.
  - 14:   Calculate  $y_{t+h}^{(l)}(\mathbf{y}_{t-1})$  by using the previous observations  $(\mathbf{y}_{t-1}, y_t^{(l)}(\mathbf{y}_{t-1}), \dots, y_{t+h-1}^{(l)}(\mathbf{y}_{t-1})$  and the structural error  $e_{t+h}$  obtained from Steps 3-12.
  - 15: **end for**
  - 16: Calculate  $y_{t+h}^{(l)}(\delta_j, \mathbf{y}_{t-1}) - y_{t+h}^{(l)}(\mathbf{y}_{t-1})$ .
  - 17: Repeat Steps 2-16  $R$  times and calculate the sample mean of  $y_{t+h}^{(l)}(\delta_j, \mathbf{y}_{t-1}) - y_{t+h}^{(l)}(\mathbf{y}_{t-1})$  over the Monte Carl repetitions  $l = 1, \dots, R$  to obtain the (counterfactual) GIRF related to the history  $\mathbf{y}_{t-1}$  for the  $j$ th structural shock of sign and size  $\delta_j$ .
- 

p = 1, M = 2, d = 2, #parameters = 19,

Regime 1

Weight param: 0.60

Regime means: 0.22, 0.87

	Y	phi0	A1		Omega	1/2	
1	y1 =	[ 0.00 ]	+	[ 0.20 0.20 ]	y1.1 +	[ 1.00 0.10 ]	eps1
2	y2	[ 1.00 ]		[ 0.20 -0.20 ]	y2.1	[ 0.10 1.00 ]	eps2

Regime 2

Weight param: 0.40

Regime means: 0.73, 1.71

	Y	phi0	A1		Omega	1/2	
1	y1 =	[ 0.00 ]	+	[ 0.30 0.30 ]	y1.1 +	[ 4.00 0.40 ]	eps1
2	y2	[ 2.00 ]		[ 0.30 -0.30 ]	y2.1	[ 0.40 4.00 ]	eps2

It is possible to include data in the models built with **STVAR** by either providing the data in the argument **data**. When the model is supplied with data, the transition weights and other data dependent statistics are calculated for the model as well. It is also possible to build models that do not satisfy the usual stability condition for all the regimes. In this case, you should set the



argument `allow_unstab` to `FALSE`.

## 11. Simulation and forecasting

### 11.1. Simulation

**sstvars** implements the S3 method `simulate` for simulating observations from STVAR processes (see `?simulate.stvar`). The method requires the process to be given as a class `stvar` object, which are typically created either by estimating a model with the function `fitSTVAR` (or `fitSSTVAR`) or by specifying the parameter values by hand and building the model with the constructor function `STVAR`. The initial values required to simulate the first  $p$  observations can be either set by hand (with the argument `init_values`) or drawn from (the stationary distribution of) some regime (with the argument `init_regime`). The argument `nsim` sets the length of the sample path to be simulated.

To give an example, the following code sets the random number generator seed to one and simulates 500 observations long sample from the STVAR model built in Section 10, drawing initial values from the first regime:

```
R> mysim <- simulate(mod122, nsim=500, init_regime=1, seed=1)
```

Our implementation of `simulate` returns a list containing the simulated sample path in `$sample`, the mixture component that generated each observation in `$component`, and the transition weights in `$transition_weights`.

### 11.2. Simulation based forecasting

Deriving multiple-steps-ahead point predictions and prediction intervals analytically for the STVAR models is complicated, so **sstvars** employs the following simulation-based method. By using the last  $p$  observations of the data up to the date of forecasting as initial values, a large number of sample paths for the future values of the process are simulated. Then, sample quantiles from the simulated sample paths are calculated to obtain prediction intervals, and the median or mean is used for point predictions. A similar procedure is also applied to forecast future values of the transition weights, which might be of interest because the researcher can often associate statistical characteristics or economic interpretations to the regimes.

Forecasting is most conveniently done with the `predict` method (see `?predict.stvar`). The available arguments include the number of steps ahead to be predicted (`nsteps`), the number sample paths the forecast is based on (`nsim`), possibly multiple confidence levels for prediction intervals (`pi`), prediction type (`pred_type`), and prediction interval type (`pi_type`). The prediction type can be either `median`, `mean` for the point forecast.

To exemplify, the following code forecasts the two-dimensional time-series of U.S. GDP and GDP deflator growth using the logistic Student's  $t$  STVAR(1,2) model `fit12` estimated in Section 4.5. The forecast is 10 steps (quarters in this case) ahead, based on 10000 Monte Carlo repetitions with the point forecast based on the mean of those repetitions. The prediction intervals are two-sided with confidence levels 0.95 and 0.90.

```
R> mypred <- predict(fit12, nsteps=10, nsim=10000, pred_type="mean",
+   pi=c(0.95, 0.90))
```

The results can be printed with the `print` method using the command `print(mypred)` or plotted with the `plot` method using the command `plot(mypred)`.



## 12. Summary

Smooth transition vector autoregressive models are a valuable tool in modelling multivariate time series in which the data generating dynamics vary in time, exhibiting gradual shifts in the autoregressive coefficients or conditional covariance matrices. We described the R package **sstvars**, which accommodates STVAR models with various transition weight functions, including exogenous weights, logistic weights (Anderson and Vahid 1998), multinomial logit weights, exponential weights (e.g., Hubrich and Teräsvirta 2013), threshold weights (Tsay 1998), and transition weights that defined as weighted relative likelihoods of the regimes corresponding to the preceding  $p$  observations (Lanne and Virolainen 2025). Currently, the accommodated conditional distributions include Gaussian distribution, Student's  $t$  distribution, Student's  $t$  distribution with mutually independent components, and skewed  $t$  distribution with mutually independent components (see Hansen 1994). The accommodated identification methods include recursive identification, identification by heteroskedasticity, and identification by non-Gaussianity. We discussed the various model specifications and several features implemented in **sstvars** for STVAR modeling: unconstrained and constrained maximum likelihood estimation of the model parameters, impulse response analysis, historical decompositions, counterfactual analysis residual based diagnostics, hypothesis testing, simulation, forecasting, and more. For convenience, we have collected some useful functions in **sstvars** to Table 1. For all the exported functions and their usage, see the reference manual.

## A table of some useful functions

Related to	Name	Description
Estimation	<code>fitSTVAR</code>	Estimate STVAR models.
	<code>fitSSTVAR</code>	Estimate or construct structural STVAR models.
	<code>alt_stvar</code>	Construct a STVAR model based on any estimation round.
	<code>filter_estimates</code>	Browse estimates, not deemed inappropriate, from various estimation rounds.
	<code>iterate_more</code>	Run more iterations of the variable metric algorithm for a preliminary estimated STVAR model.
Estimates	<code>print (method)</code>	Print the estimates or their approximate standard errors.
	<code>summary (method)</code>	Detailed printout of the model.
	<code>plot (method)</code>	Plot the series with the fitted transition weights of the model.
	<code>get_foc</code>	Calculate numerically approximated gradient of the log-likelihood function evaluated at the estimate.
	<code>get_soc</code>	Calculate eigenvalues of numerically approximated Hessian of the log-likelihood function evaluated at the estimate.
Diagnostics	<code>profile_logliks</code>	Plot the graphs of the profile log-likelihood functions about the estimate.
	<code>profile_struct_shocks</code>	Plot the time series of the structural shocks.
	<code>Portmanteau_test</code>	Calculate the (adjusted) Portmanteau test.
	<code>diagnostic_plot</code>	Plot residual diagnostics (raw or standardized residuals).
	<code>predict (method)</code>	Forecast future observations and transition weights of the process.
Simulation	<code>simulate (method)</code>	Simulate from a STVAR process.
Create model	<code>STVAR</code>	Construct a STVAR model based on given parameter values.
Hypothesis testing	<code>LR_test</code>	Calculate likelihood ratio test.
	<code>Wald_test</code>	Calculate Wald test.
	<code>Rao_test</code>	Calculate Rao's test.
Impulse responses	<code>GIRF</code>	Estimate generalized impulse response functions.
	<code>GFEVD</code>	Estimate generalized forecast error variance decomposition.
	<code>linear_IRF</code>	Estimate linear impulse response functions.
Hist. decomp.	<code>hist_decomp</code>	Compute historical decompositions.
Counterfactuals	<code>cfact_hist</code>	Simulate historical counterfactuals.
	<code>cfact_fore</code>	Simulate counterfactual forecast scenarios.
	<code>cfact_girf</code>	Simulate counterfactual generalized impulse response functions.
Other	<code>bound_JSR</code>	Calculate bounds for the joint spectral radius of the companion form AR matrices of the regimes.
	<code>swap_parametrization</code>	Swap between mean and intercept parametrizations
	<code>uncond_moments</code>	Calculate unconditional moments of the regimes.
	<code>check_params</code>	Check whether given parameter values satisfy our assumptions.
	<code>swap_B_signs</code>	Swap the signs of the columns of the impact matrix of models identified by heteroskedasticity or non-Gaussianity.
	<code>reorder_B_columns</code>	Reorder the columns of the impact matrix of models identified by heteroskedasticity or non-Gaussianity.

Table 1: Some useful functions in **sstvars** sorted according to their usage. The note "method" in parentheses after the name of a function signifies that it is an S3 method for a class **stvar** object (often generated by the function `fitSTVAR`, `fitSSTVAR` or `STVAR`).

## References

- Anderson H, Vahid F (1998). “Testing multiple equation systems for common nonlinear components.” *Journal of Econometrics*, **84**(1), 1–36. doi:[10.1016/S0304-4076\(97\)00076-6](https://doi.org/10.1016/S0304-4076(97)00076-6).
- Bernanke B, Gertler M, Watson M (1997). “Systematic monetary policy and the effects of oil price shocks.” *Brookings Papers on Economic Activity*, **1**, 91—142.
- Burgard J, Neuenkirch M, Nöckel M (2019). “State-Dependent Transmission of Monetary Policy in the Euro Area.” *Journal of Money, Credit and Banking*, **51**(7), 2053–2070. doi:[10.1111/jmcb.12592](https://doi.org/10.1111/jmcb.12592).
- Buse A (1982). “The Likelihood Ratio, Wald, and Lagrange Multiplier Tests: An Expository Note.” *The American Statistician*, **36**(3a), 153–157. doi:[10.1080/00031305.1982.10482817](https://doi.org/10.1080/00031305.1982.10482817).
- Carriero A, Marcellino M, Tornese T (2024). “Blended identification in structural VARs.” *Journal of Monetary Economics*, **146**, Article 103581. doi:[10.1016/j.jmoneco.2024.103581](https://doi.org/10.1016/j.jmoneco.2024.103581).
- Chang CT, Blondel V (2013). “An experimental study of approximation algorithms for the joint spectral radius.” *Numerical Algorithms*, **64**, 181–202. doi:[10.1007/s11075-012-9661-z](https://doi.org/10.1007/s11075-012-9661-z).
- Gripenberg G (1996). “Computing the joint spectral radius.” *Linear algebra and its applications*, **234**, 43–60. doi:[10.1016/0024-3795\(94\)00082-4](https://doi.org/10.1016/0024-3795(94)00082-4).
- Hansen BE (1994). “Autoregressive Conditional Density Estimation.” *International Economic Review*, **35**(3), 705–730.
- Herwartz H, Lütkepohl H (2014). “Structural vector autoregressions with Markov switching: Combining conventional with statistical identification of shocks.” *Journal of Econometrics*, **183**(1), 104–116. doi:[doi.org/10.1016/j.jeconom.2014.06.012](https://doi.org/10.1016/j.jeconom.2014.06.012).
- Hubrich K, Teräsvirta T (2013). “Thresholds and Smooth Transitions in Vector Autoregressive Models.” *CREATES Research Paper 2013-18*, Aarhus University. URL [https://pure.au.dk/ws/files/54473123/rp13\\_18.pdf](https://pure.au.dk/ws/files/54473123/rp13_18.pdf).
- Jungers R (2023). *The JSR toolbox* (<https://www.mathworks.com/matlabcentral/fileexchange/33202-the-jsr-toolbox>), MATLAB Central File Exchange.
- Kalliovirta L, Meitz M, Saikkonen P (2016). “Gaussian mixture vector autoregression.” *Journal of Econometrics*, **192**(2), 465–498. doi:[10.1016/j.jeconom.2016.02.012](https://doi.org/10.1016/j.jeconom.2016.02.012).
- Kheifets I, Saikkonen P (2020). “Stationarity and ergodicity of vector STAR models.” *Econometric Reviews*, **39**(407–414), 1311–1324. doi:[10.1080/07474938.2019.1651489](https://doi.org/10.1080/07474938.2019.1651489).
- Kilian L, Lewis L (2011). “Does the Fed Respond to Oil Price Shocks?” *The Economic Journal*, **121**(555). doi:[10.1111/j.1468-0297.2011.02437.x](https://doi.org/10.1111/j.1468-0297.2011.02437.x).
- Kilian L, Lütkepohl H (2017). *Structural Vector Autoregressive Analysis*. 1st edition. Cambridge University Press, Cambridge. doi:[10.1017/9781108164818](https://doi.org/10.1017/9781108164818).
- Koop G, Pesaran M, Potter S (1996). “Impulse response analysis in nonlinear multivariate models.” *Journal of Econometrics*, **74**(1), 119–147. doi:[10.1016/0304-4076\(95\)01753-4](https://doi.org/10.1016/0304-4076(95)01753-4).
- Lanne M, Meitz M, Saikkonen P (2017). “Identification and estimation of non-Gaussian structural vector autoregressions.” *Journal of Econometrics*, **196**(2), 288–304. doi:[j.jeconom.2016.06.002](https://doi.org/10.1016/j.jeconom.2016.06.002).

- Lanne M, Nyberg H (2016). “Generalized Forecast Error Variance Decomposition for Linear and Nonlinear Multivariate Models.” *Oxford Bulletin of Economics and Statistics*, **78**(4), 595–603. doi:[10.1111/obes.12125](https://doi.org/10.1111/obes.12125).
- Lanne M, Virolainen S (2025). “A Gaussian smooth transition vector autoregressive model: An application to the macroeconomic effects of severe weather shocks.” *Unpublished working paper, available as arXiv:2403.14216*. doi:[10.48550/arXiv.2403.14216](https://doi.org/10.48550/arXiv.2403.14216). URL <https://doi.org/10.48550/arXiv.2403.14216>.
- Lütkepohl H (2005). *New Introduction to Multiple Time Series Analysis*. 1st edition. Springer, Berlin. doi:[10.1007/978-3-540-27752-1](https://doi.org/10.1007/978-3-540-27752-1).
- Lütkepohl H, Netsunajev A (2017). “Structural vector autoregressions with smooth transitions in variances.” *Journal of Economic Dynamics & Control*, **84**, 43–57. doi:[10.1016/j.jedc.2017.09.001](https://doi.org/10.1016/j.jedc.2017.09.001).
- Meitz M, Preve D, Saikkonen P (2023). “A mixture autoregressive model based on Student’s  $t$ -distribution.” *Communications in Statistics - Theory and Methods*, **52**(2), 499–515. doi:[10.1080/03610926.2021.1916531](https://doi.org/10.1080/03610926.2021.1916531).
- Muirhead R (1982). *Aspects of Multivariate Statistical Theory*. 1st edition. John Wiley & Sons, Hoboken, New Jersey. doi:[10.1002/9780470316559](https://doi.org/10.1002/9780470316559).
- Nash J (1990). *Compact Numerical Methods for Computers. Linear Algebra and Function Minimization*. 2nd edition. Adam Hilger, Bristol and New York. doi:[10.1201/9781315139784](https://doi.org/10.1201/9781315139784).
- R Core Team (2022). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Rigobon R (2003). “Identification through Heteroskedasticity.” *The Review of Economics and Statistics*, **85**(4), 777–792. doi:[10.1162/003465303772815727](https://doi.org/10.1162/003465303772815727).
- Saikkonen P (2008). “Stability of Regime Switching Error Correction Models under Linear Cointegration.” *Econometric Theory*, **24**(1), 294–318. doi:[10.1017/S0266466608080122](https://doi.org/10.1017/S0266466608080122).
- Solymos P, Zawadzki Z (2020). *pbapply: Adding Progress Bar to \*apply\* Functions*. R package version 1.4-3, URL <https://CRAN.R-project.org/package=pbapply>.
- Teräsvirta T (1994). “Specification, Estimation, and Evaluation of Smooth Transition Autoregressive Models.” *Journal of the American Statistical Association*, **89**(425), 208–2187. doi:[10.2307/2291217](https://doi.org/10.2307/2291217).
- Tsay R (1998). “Testing and Modeling Multivariate Threshold Models.” *Journal of the American Statistical Association*, **93**(443), 1188–1202. doi:[10.1080/01621459.1998.10473779](https://doi.org/10.1080/01621459.1998.10473779).
- Virolainen (2025a). “Identification by non-Gaussianity in structural threshold and smooth transition vector autoregressive models.” *Unpublished working paper, available as arXiv:2404.19707*.
- Virolainen S (2018a). *gmvarKit: Estimate Gaussian and Student’s  $t$  Mixture Vector Autoregressive Models*. R package version 2.1.0 available at CRAN: <https://CRAN.R-project.org/package=gmvarKit>, URL <https://CRAN.R-project.org/package=gmvarKit>.
- Virolainen S (2018b). *uGMAR: Estimate Univariate Gaussian and Student’s  $t$  Mixture Autoregressive Models*. R package version 3.4.2 available at CRAN: <https://CRAN.R-project.org/package=uGMAR>, URL <https://CRAN.R-project.org/package=uGMAR>.

- Virolainen S (2022a). *Essays on mixture autoregressive models with applications to macroeconomics and finance*. Ph.D. thesis, University of Helsinki. URL <http://hdl.handle.net/10138/350319>.
- Virolainen S (2022b). “A mixture autoregressive model based on Gaussian and Student’s *t*-distributions.” *Studies in Nonlinear Dynamics & Econometrics*, **26**(4), 559–580. doi:10.1515/snde-2020-0060.
- Virolainen S (2025b). “A statistically identified structural vector autoregression with endogeneously switching volatility regime.” *Journal of Business & Economic Statistics*, **43**(1), 44–54. doi:10.1080/07350015.2024.2322090.
- Wong B (2018). “Historical Decompositions for Nonlinear Vector Autoregression Models.” *CAMA Working Paper No. 62/2017*, available as SSRN:3057759. doi:10.2139/ssrn.3057759.

**Affiliation:**

Savi Virolainen  
Faculty of Social Sciences  
University of Helsinki  
P. O. Box 17, FI-0014 University of Helsinki, Finland  
E-mail: [savi.virolainen@helsinki.fi](mailto:savi.virolainen@helsinki.fi)